

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(51) Publication number:

**0 362 179  
A3**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **89870129.7**

(22) Date of filing: **24.08.89**

(51) Int. Cl.<sup>5</sup> **C12N 1/18, C12N 15/81,  
C12N 15/62, C12P 21/02,  
C12N 15/49, C12N 15/25,  
C12N 15/31, C07K 15/00,  
C12N 15/19, A61K 39/10,  
//(C12N1/18,C12R1:865)**

(70) Priority: **25.08.88 US 236699**

(43) Date of publication of application:  
**04.04.90 Bulletin 90/14**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(88) Date of deferred publication of the search report:  
**11.07.90 Bulletin 90/28**

(71) Applicant: **SMITHKLINE BEECHAM  
CORPORATION  
One Franklin Plaza  
Philadelphia Pennsylvania 19103(US)**

(72) Inventor: **Bruck, Claudine  
Avenue du Gris Moulin, 53  
B-1310 La Hulpe(BE)  
Inventor: Cohen, Joseph  
Rue Edmond Picard, 52  
B-1180 Bruxelles(BE)  
Inventor: Gorman, Jessica Angel  
152 Ridgfield Road  
Newtown Square Pennsylvania 19073(US)  
Inventor: Koltin, Yigal  
16B Hashalom Street  
Hod Hasharon(IL)  
Inventor: Locht, Camille  
Avenue Bel Horizon, 40  
B-1320 Genval (Rixensart)(BE)  
Inventor: Van Wijnendaele, Frans  
Teckerstraat, 12  
B-3052 Ottenburg(BE)**

(74) Representative: **Giddings, Peter John, Dr.  
SmithKline Beecham, Corporate Patents,  
Mundells  
Welwyn Garden City, Hertfordshire AL7  
1EY(GB)**

**0 362 179 A3**

(54) **Recombinant saccharomyces.**

secretory heterologous gene products which are under control of a signal sequence derived from Candida or Saccharomyces. Also provided are a method for characterizing the function of the Candida DNA se-



European Patent  
Office

**PARTIAL EUROPEAN SEARCH REPORT**  
which under Rule 45 of the European Patent Convention  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

Application number

EP 89 87 0129

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	EP-A-0 173 668 (SMITHKLINE Beckman Corp.)  * Page 15, lines 9-29 *  --	1-20	C 12 N 1/18 C 12 N 15/81 C 12 N 15/62 C 12 P 21/02 C 12 N 15/49 C 12 N 15/25 C 12 B 15/31 C 07 K 15/00 C 12 N 15/19 A 61 K 39/10// (C 12 N 1/18, C 12 R 1:865)
A	SCIENCE, vol. 229, September 20, 1985, pages 1219-1224 Washington, DC, US; R.A. SMITH et al.: "Heterologous protein secretion from yeast"  --		
X	VACCINE, vol. 5, no. 2, June 1987, pages 90-101, Butterworth & Co. Publ. Ltd., Guildford, GB; P.J. BARR et al.: "Antigenicity and immunogenicity of domains of the human immunodeficiency virus (HIV) envelope polypeptide expressed in the yeast <i>Saccharomyces cerevisiae</i> "  * Figure 2 * -----	22-24	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 12 N C 12 P
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-32, 34-39 Claims searched incompletely: 33 Claims not searched: 33 Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (see Art. 52(4) of the European Patent Convention).</p>			
Place of search Munich		Date of completion of the search 10-12-1989	Examiner VAN PUTTEN

EPO Form 1101

LEGEND

X : particularly relevant if taken alone  
Y : particularly relevant if combined with another document of the same category  
A : technological background  
O : non-written disclosure  
P : intermediate document

Legend (continued)

... of principle under which the ...  
earlier patent document, but published ...  
after the filing date  
... document cited in the application  
... document cited for other reasons  
... member of the same patent family, corresponding document



European Patent  
Office

### CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

### X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

See annex -B-

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims.

namely claims: 1-16, 22-24 and 17-20 partially



# LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions,

namely:

1. Claims 1-16, 22-24 and 17-20 partially:  
A transformed host microorganism of the genus Saccharomyces which comprises a functional DNA sequence derived from the same or substantially the same coding sequence as the signal sequence of the C.albicans glucoamylase gene.  
A recombinant DNA plasmid comprising a functional DNA sequence containing a coding sequence the same or substantially the same as the signal sequence of the C. albicans glucoamylase gene.  
A method of expressing a functional DNA sequence comprising a sequence the same or substantially the same as the signal sequence of the C. albicans glucoamylase gene which comprises transforming a host microorganism of the genus Saccharomyces with the functional DNA sequence, and culturing the transformed host under suitable conditions such that the functional DNA sequence is expressed.  
An HIV-1 glycoprotein gp 160.
2. Claims 21 and 17-20 partially:  
A method of expressing a functional DNA sequence comprising a sequence the same or substantially the same as the promoter sequence of the C. albicans glucoamylase gene which comprises transforming a host microorganism of the genus Saccharomyces with the functional DNA sequence, and culturing the transformed host under suitable conditions such that the functional DNA sequence is expressed.  
A C.albicans glucoamylase gene promoter sequence.
3. Claim 25: TGF-alpha
4. Claim 26-28,31,32,34-39:  
S1-GA protein  
Vaccin comprising this protein  
a recombinant DNA molecule  
A synthetic DNA molecule  
Host cell transformed with this DNA
5. Claim 29: IL1-beta
6. Claim 30: IL1-alpha

## Ginger Dreger

---

**From:** Carol Peterson  
**Sent:** Thursday, June 21, 2001 1:50 PM  
**To:** Ginger Dreger  
**Subject:** RE: GENHE.000GEN

Hi! I put the results on file:h:\docs\csp\wax.doc. I tried various combinations of all of the terms, but all of the hits look like duds to me. I think that it might be because this technique is used in other pursuits so is mentioned somewhere in the full text but is not the main point of the patents. I put lots of notes in the margins by the statements from which I printed, and after awhile I just printed titles (instead of including abstracts) so as to cut down the cost and verbage since they may be irrelevant. If you see anything that interests you please let me know and I will get the abstract. I also looked for the possible assignees which you mentioned, and the only one that looked good was the US 6183995--the one that you already have. It did NOT turn up in the searches with the terms so I think that we need to modify the strategy. I will give you a call also to discuss.

-----Original Message-----

**From:** Ginger Dreger  
**Sent:** Thursday, June 21, 2001 1:01 PM  
**To:** Carol Peterson  
**Subject:** RE: GENHE.000GEN

Tha patent is 6,183,995 - sorry for the typo!!!

-----Original Message-----

**From:** Carol Peterson  
**Sent:** Thursday, June 21, 2001 11:51 AM  
**To:** Ginger Dreger  
**Subject:** RE: GENHE.000GEN

I pulled up this patent number and it's a wrong #-some sort of gasket. Carlos looked briefly through a stack of patents on your desk, but there was nothing close. I am going to go ahead using the terms that you've given and then go back and read that patent when I get the #.

-----Original Message-----

**From:** Ginger Dreger  
**Sent:** Thursday, June 21, 2001 11:30 AM  
**To:** Carol Peterson  
**Subject:** RE: GENHE.000GEN

Great!

-----Original Message-----

**From:** Carol Peterson  
**Sent:** Thursday, June 21, 2001 11:28 AM  
**To:** Ginger Dreger  
**Cc:** Suzi Kortenkamp  
**Subject:** RE: GENHE.000GEN

Hi! Sorry to have been incommunicado on this matter. I was planning to do it today and things got rather hectic. I am starting on it right now. I'll give you a call too Ginger.

-----Original Message-----

**From:** Ginger Dreger  
**Sent:** Wednesday, June 20, 2001 2:41 PM  
**To:** Carol Peterson  
**Subject:** FW: GENHE.000GEN

Hi Carol,

-----Original Message-----

**From:** Ginger Dreger  
**Sent:** Monday, June 18, 2001 2:22 PM

To:  
Subject:

Library  
GENHE.000GEN

**PRIVILEGED AND CONFIDENTIAL COMMUNICATION**

The information contained in this electronic transmission, and any documents attached hereto, contain confidential information that is legally privileged and confidential. The information is intended only for the use of the recipient named above. If you have received this electronic message in error, please notify the sender and delete the electronic message; and any disclosure, copying, distribution or the taking of any action in reliance on the contents of this information is strictly prohibited.

As usual, I need your urgent help with a patent search project. On behalf of a new client, Genomic Health, we were asked to urgently search for patents and published patent applications worldwide with CLAIMS that might cover measuring gene expression using RNA isolated from wax-embedded tissue specimens.

Samples taken from solid tumors, e.g. by biopsy, are usually preserved as paraffin-embedded, formalin-fixed tissues. From these tissues, RNA can be subsequently isolated, and gene expression within the tissue analyzed, e.g. by reverse transcription PCR (RT-PCR). I suggest that you search for the appearance of the following terms in the claims:

(paraffin-embedded or wax-embedded) and RNA and expression optionally coupled with RT-PCR and/pr tumor.

US 6,183,905 is right on the point, i.e. it might be helpful in designing the search.

Potential applicants/assignees: LifeSpan BioSciences, Inc., Exact Laboratories, Inc., or Exact Sciences, or Response Genetics.

Once we have the results of the patent search, we will need to conduct a prior art search, and provide an analysis all within two weeks. As a result, this is very urgent.

Please contact me if you need any further information. Otherwise, please send me the results, including abstracts/claims of the patent publications found.

I appreciate your help!

Ginger



HOME
Office Services
Attorney Profiles
Online Manuals
Departmental Areas
Document Search Page
Staff Evaluation
Employee Telephone List
File Locations

	Thursday, June 21, 2001
<b>KMOB NEWS</b>  Web Links  Holiday Party  National Library Week	<b>WORLD NEWS</b>



(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 362 179  
A2**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 89870129.7

(22) Date of filing: 24.08.89

(51) Int. Cl.<sup>5</sup>: **C12N 1/18 , C12N 15/81 ,  
C12N 15/62 , C12P 21/02 ,  
C12N 15/49 , C12N 15/25 ,  
C12N 15/31 , C07K 15/00 ,  
C12N 15/19 , A61K 39/10 ,  
/(C12N1/18,C12R1:865)**

(30) Priority: 25.08.88 US 236699

(43) Date of publication of application:  
04.04.90 Bulletin 90/14

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(71) Applicant: **SMITHKLINE BEECHAM  
CORPORATION**  
One Franklin Plaza  
Philadelphia Pennsylvania 19103(US)

(72) Inventor: **Bruck, Claudine**  
Avenue du Gris Moulin, 53  
B-1310 La Hulpe(BE)  
Inventor: **Cohen, Joseph**  
Rue Edmond Picard, 52  
B-1180 Bruxelles(BE)  
Inventor: **Gorman, Jessica Angel**  
152 Ridgefield Road  
Newtown Square Pennsylvania 19073(US)  
Inventor: **Koltin, Yigal**  
16B Hashalom Street  
Hod Hasharon(IL)  
Inventor: **Locht, Camille**  
Avenue Bel Horizon, 40  
B-1320 Genval (Rixensart)(BE)  
Inventor: **Van Wijnendaele, Frans**  
Teckerstraat, 12  
B-3052 Ottenburg(BE)

(74) Representative: **Tasset, Gérard**  
Smithkline Biologicals s.a. Rue de l'Institut  
89  
B-1330 Rixensart(BE)

EP 0 362 179 A2

(54) Recombinant saccharomyces.

(57) The invention relates to a method for cloning and expressing DNA sequences derived from Candida in Saccharomyces; the method consists in inserting a DNA sequence derived from Candida into a plasmid vector, transforming a Saccharomyces cell with the recombinant plasmid, and selecting the transformed cells. The invention also relates to a recombinant plasmid useful for performing such transformation.

## RECOMBINANT SACCHAROMYCES

This invention is related generally to cloning and expression of DNA sequences from Candida in Saccharomyces. More particularly, the invention relates to the expression of Candida gene sequences in Saccharomyces cells to complement Saccharomyces host cell function and to the expression and secretion of heterologous genes under the control of Candida promoter/signal sequences in Saccharomyces cells.

5 This invention also relates to a vaccine against whooping cough which comprises a truncated B. pertussis S1 subunit protein expressed and secreted in a Saccharomyces cell under the control of Candida promoter/signal sequences.

10

Summary of the Invention

One aspect of this invention is a transformed host microorganism of the genus Saccharomyces which comprises a functional DNA sequence derived from a variety of Candida species.

15 Another aspect of this invention is a recombinant plasmid comprising a replicator region and a functional DNA sequence derived from Candida.

This invention also provides a promoter sequence and a yeast secretory signal sequence derived from the C. albicans glucoamylase gene for use in expression systems in yeast.

20 Still another aspect of this invention provides a recombinant plasmid comprising a yeast promoter, the signal sequence from the C. albicans glucoamylase gene and a heterologous structural gene, capable upon transformation and expression in Saccharomyces of producing a hybrid fused gene product.

This invention also relates to a method of expressing a functional DNA sequence derived from Candida which comprises transforming a host microorganism of the genus Saccharomyces with the functional DNA sequence, and culturing the transformed host under suitable conditions such that the functional DNA

25 sequence is expressed.

Yet another aspect of this invention is a method of characterizing a DNA sequence derived from Candida by its function which comprises (a) transforming a host microorganism of the genus Saccharomyces with the DNA sequence, wherein said host microorganism lacks a known function, so that the Candida DNA can complement the host microorganism; and (b) assaying for presence of the function in the

30 transformed host microorganism.

This invention also relates to a method of producing a heterologous polypeptide in Saccharomyces which comprises culturing a transformed host microorganism of the genus Saccharomyces in suitable culturing media wherein said host microorganism comprises a promoter and/or signal sequence derived from Candida fused, directly or indirectly, in phase and in proper orientation to a heterologous structural

35 gene which codes for the polypeptide. Where the signal sequence is used in the method, the polypeptide is expressed and secreted from the yeast cell host.

Thus, a further aspect of this invention includes heterologous polypeptides or fragments thereof secreted from Saccharomyces host cells according to this invention. Among these secreted polypeptides is a recombinant HIV-T glycoprotein gp160 unprocessed into its gp120 and gp41 components.

40 The present invention also relates to a vaccine for stimulating protection against whooping cough, wherein such vaccine comprises an immunoprotective and non-toxic amount of S1-GA protein. Such vaccine may comprise such protein alone or in conjunction with other antigens and/or adjuvants. In addition, the present invention also provides a method for inoculating a human against whooping cough which comprises administering the vaccine of this invention to such human.

45

Brief Description of the Drawings

Fig. 2 depicts the expression vector employed in Example 1.

Fig. 3 presents the nucleotide and amino acid sequence of the Sig A subunit of the C. albicans glucoamylase gene.

Fig. 4 presents the nucleotide and amino acid sequence of the synthetic linker employed in Example

11.

Fig. 5 presents the DNA and amino acid sequence of TGF- $\alpha$ .

#### Detailed Description of the Invention

5

The transformed host microorganism of the subject invention is any member of the genus Saccharomyces into which a functional DNA sequence derived from Candida has been introduced. Saccharomyces cerevisiae is preferred because it is the most well studied member of the genus and has been the species most frequently employed for recombinant DNA purposes.

The recombinant plasmid of the subject invention comprised a functional DNA sequence derived from any species of the genus Candida including species, Candida albicans, Candida tropicalis, Candida stellatoidea, Candida pseudotropicalis, Candida parapsilosis, Candida quillieromondi, Candida utilis and Candida kruisi. DNA derived from Candida albicans is most preferred because it was the first DNA utilized in the subject invention since it is derived from the genus member most likely to cause candidiasis.

By the term "functional DNA sequence" is meant any discrete region of DNA derived directly or indirectly from Candida which functions in Candida as a complete gene expression unit, a structural gene, a promoter, a signal sequence or a regulatory region. By "complete gene expression unit" is meant a structural gene and the promoter and regulatory regions required for its transcription and translation. By "promoter" is meant any region upstream of a structural gene which permits binding of RNA polymerase and transcription to occur. By "regulatory region" is meant any region which regulates transcription of the gene. By "structural gene" is meant a coding sequence for a polypeptide which serves to be the template for the synthesis of mRNA. By "signal sequence" is meant a region upstream of a structural gene which targets the struction gene to the secretion pathway of the host.

The recombinant plasmid of the subject invention also comprises a replicator region, preferably a Saccharomyces or Candida replicator region. By "replicator region" is meant any DNA fragment, preferably derived directly or indirectly from yeast, such as Saccharomyces or Candida, which can be used to maintain, extrachromosomally, a functional DNA sequence derived from Candida in the host Saccharomyces microorganism. Particularly preferred plasmids are those comprising a Saccharomyces-Escherichia coli shuttle vector such as the one described by Broach et al. Gene, 8:121-133 (1979). Illustrative of vectors which can be used in this invention by cloning therein a functional DNA sequence derived from Candida are the following plasmids which are publicly available, e.g., from the American Type Culture Collection, Rockville, Maryland, U.S.A. (ATCC accession number in parentheses): pBIT-1 (37087), pBTI-7 (37088), pBTI-9 (37089), pBTI-10 (37090), pLC544 (37013), pRB3 (37059), pRB4 (37058), pRB5 (37051), pRB8 (37109), pRB9 (37057), pRB10 (37056), pRB18 (37092), pRB20 (37054), pRB21 (37055), pRB22 (37053), pRB12 (37061), pRB14 (37063), pRB15 (37062), pRB16 (37060), pRB19 (37064), YEp2 (37076), YRp17 (37078), YEp13 (37115).

The Candida DNA sequence can be prepared by known techniques such as the one described by Rigsby et al., Molec. Cell. Biol., 2:853-862 (1982). The isolated Candida DNA can then be further fragmented by treatment with one or more restriction enzymes.

A recombinant plasmid containing a functional Candida DNA sequence and containing an Escherichia coli replicator region can be used to transform Escherichia coli to amplify the plasmid. Such transformation and amplification can be accomplished by known techniques. The recombinant plasmid is then retrieved from E. coli by known techniques.

The retrieved recombinant plasmid can then be used to transform a host microorganism belonging to the genus Saccharomyces. Such transformation can be accomplished by the random insertion of recombinant plasmid DNA fragments into covalently closed circular yeast plasmids and (a) transforming spheroplasts therewith, such as by the method described by Hinnen et al., Proc Nat'l. Acad Sci. USA, 75:1929-1933 (1978); or (b) by using the lithium acetate or lithium chloride method of yeast transformation such as described by Ito et al., J. Bacteriol., 153:163(1983). DNA from Candida can also be transformed into a Saccharomyces microorganism by fusion of Candida and Saccharomyces protoplasts. The introduced Candida DNA can be integrated into the host Saccharomyces microorganism's chromosomal DNA, such as by recombination or can remain extrachromosomal. Thus, yet another exemplary method is transformation

of a Candida microorganism with Candida DNA which optionally has flanking sequences homologous to sequences in the host Candida microorganism's chromosomal DNA.

Promoter regions and signal sequences derived from Candida by the process of the subject invention can be used to express coding sequences from Candida, Saccharomyces or other organisms. Uses of

cells in the transformed Saccharomyces host. Such promoters and/or signal sequences can be fused to a heterologous structural gene, such as sequences coding for insulin, interferon, growth hormone bacterial and viral proteins, etc., directly by fusing all or a portion of the promoter's and/or signal sequence's natural nucleotide sequence to the coding sequence for the desired polypeptide, thereby producing a hybrid gene. For example, one such promoter and signal region is derived from C. albicans glucoamylase gene. This functional DNA sequence when inserted in a plasmid and fused to a heterologous structural gene enables the expression and secretion of a hybrid fusion protein. An alternative expression system employs another yeast promoter and the C. albicans glucoamylase gene signal sequence for expression and secretion of a desired fusion gene. When Saccharomyces host microorganisms, which have been transformed by a plasmid containing such hybrid gene, are cultured in suitable culturing media they produce the polypeptide. Under control of the appropriate promoter and signal sequence, the polypeptide may be secreted into the culture medium.

In addition to those Candida promoters disclosed herein, additional promoters can be discovered by cloning random fragments of Candida DNA in Saccharomyces plasmids designed for promoter selection. 15 Such plasmids can be designed by known techniques such as the one described by Casadaban et al., Methods in Enzymology, 100: 293-308(1983).

Structural genes derived from Candida can be expressed in Saccharomyces from their natural Candida promoters, from heterologous promoters derived from Candida or Saccharomyces, or from other functional heterologous promoters which are derived from other organisms or produced from synthetic oligonucleotides and which function in Saccharomyces. Similarly, structural genes of non-Candida origin may be expressed from a variety of promoters including the glucoamylase gene promoter with its signal sequence.

Many known Saccharomyces mutants are defective in defined functions, such as a specific step in leucine or histidine biosynthesis. This function could be supplied if such mutants were transformed with DNA sequences derived from Candida which contain genes coding for such functions. Where the function to be characterized is not already defective in an available Saccharomyces strain, such mutations can be induced by known techniques, such as by ultraviolet radiation, chemical mutagenesis, or genetic manipulation. In some cases the function displayed by the transformed host microorganism will be one which is naturally not present in Saccharomyces. For example, a host Saccharomyces cerevisiae microorganism can be transformed by a functional DNA sequence according to the method of this invention to display functions it does not naturally possess such as sorbitol utilization, maltose utilization by isomaltase catalyzed cleavage of  $\alpha$ -1-6-glucosidic bonds (such as  $\alpha$ -methyl-D-glucopyranoside hydrolysis), soluble starch utilization by glucoamylase catalyzed cleavage of  $\alpha$ -1-4-glucosidic bonds or resistance to an anti-fungal agent.

The development of a genomic library covering the entire genome of *Candida* enables the characterization of the functions of the DNA derived from *Candida*. This will permit the genetics of this asexual pathogen to be more closely analyzed than has been possible in the past and to facilitate the development of anti-candidal agents. The preferred method of characterizing the functions of DNA derived from *Candida albicans* in *Saccharomyces cerevisiae* so that the *Candida* genes can be studied in the background of a well characterized eukaryotic organism. Thus, the genetics of *Candida albicans* can be studied by isolation of genes via complementation of functions displayed by transformed *Saccharomyces cerevisiae*. The abundance of available *Saccharomyces cerevisiae* mutant strains, together with the possibility of selection of additional mutants, makes it possible to effect such complementation. Therefore, it is possible to select for *Candida albicans* genes associated with any one of a number of functions, such as those associated with defined biochemical reactions, or those involved in cell division or cell structure. The availability of the genomic library and a system for the identification of the selected genes offers a new approach to resolve some of the basic differences that make one organism economically useful and another organism a serious pathogen.

Using the method of this invention plasmids carrying one or more functional C. albicans genes which complement a Saccharomyces mutation or confer a new characteristic to the Saccharomyces strain transformed therewith can be isolated and characterized. Such functional C. albicans genes can include a gene complementing the S. cerevisiae trp1 mutation, a gene complementing the S. cerevisiae his3 mutation, a gene conferring high level resistance to the anti-fungal agent benomyl, a gene allowing for growth on soluble starch (i.e., a gene encoding amylase), a gene allowing for growth on insoluble starch (i.e., a gene encoding pullulanase), a gene allowing for growth on cellobiose (i.e., a gene encoding cellobiohydrolase) or other polysaccharides, and a gene encoding a protease.

Plasmids containing such genes can also be used to transform Saccharomyces strains deficient in the corresponding genes. Each of the promoters of the C. albicans secreted, extracellular, and soluble starch utilization genes is also useful as a regulatable promoter for a Saccharomyces microorganism transformed with a plasmid containing such a promoter. Plasmids containing C. albicans

functional DNA sequences are useful for production in Saccharomyces of heterologous gene products, such as pharmaceuticals, vaccinal antigens and hormones.

Also according to a method of this invention, a heterologous gene encoding, e.g. an HIV-1 surface glycoprotein (gp160), a B. pertussis toxin subunit (S1), Transforming Growth Factor- $\alpha$  (TGF- $\alpha$ ) and Interleukin 1 $\beta$  (IL1- $\beta$ ) may be expressed and secreted in Saccharomyces under control of a yeast promoter and the signal sequence of the C. albicans glucoamylase gene. Plasmids containing a selected yeast promoter, the glucoamylase signal sequence and the heterologous structural gene may be transformed into a selected Saccharomyces yeast cell. A gene product is expressed thereby as a hybrid or fusion protein or polypeptide and is processed and secreted into the culture medium. Secretion of the fusion protein enables purification of the protein to be greatly simplified and allows post-translational modifications of the protein, e.g. glycosylation, to be effected.

This invention also relates to a vaccine against whooping cough (pertussis). Pertussis toxin is a protein exotoxin of B. pertussis which plays a major role in the pathogenesis of whooping cough and is believed to be the major protective antigen of B. pertussis. [See, A.A. Weiss et al., *Ann. Rev. Microbiol.*, 40:661 (1986)]. Pertussis toxin is a hexameric protein composed of five dissimilar subunits called S1 through S5 according to their decreasing molecular weights. [See, M. Tamura et al., *Biochem.*, 21:5516 (1982)]. Pertussis toxin is structured in an A-B model in which the B-oligomer (subunits S2 through S5) is responsible for the binding of the toxin to its receptor in the target cell membranes, and the A-protomer (S1 subunit) contains enzymatic activity.

The nucleotide coding sequence of the entire pertussis toxin operon has been determined. See, e.g., Keith, et al., United States Patent Application Serial No. (U.S.S.N.) 843,727, filed March 26, 1986, the entire disclosure of which is hereby incorporated by reference. Furthermore DNA containing the pertussis toxin coding sequence may be isolated from any publicly available B. pertussis strain. For example, various strains of B. pertussis are publicly available from commercial depositories, e.g., from the American Type Culture Collection, Rockville, Maryland U.S.A. Such isolation may be effected by the method of Keith, et al., U.S.S.N. 843,727, supra, or any other conventional method.

The present invention is based on the discovery that S1-GA protein has the ability to elicit a protective immunogenic response. Thus, the present invention relates to S1-GA protein, the S1-GA protein coding sequence, a vaccine for stimulating protection against whooping cough, wherein such vaccine comprises an immunoprotective and non-toxic amount of S1-GA protein. Such vaccine may comprise such protein alone or in conjunction with other antigens and/or adjuvants. In addition, the present invention also provides a method for inoculating a human against whooping cough which comprises administering the vaccine of this invention to such human.

By the term 'immunoprotective amount' as used herein is meant an amount of S1-GA protein which will elicit a sufficient protective immune response against pertussis after such amount of such protein is administered to a human host.

By the term 'S1-GA protein' is meant a soluble form of the pertussis toxin (PT) S1 subunit protein which has been expressed by a host Saccharomyces cell transformed by a recombinant DNA fragment comprising 186 amino acids of the S1 subunit coding sequence (amino acids 2 to 187 of the N-terminal of the mature S1 protein) fused, in phase, to (a) the Candida albicans glucoamylase signal peptide coding sequence or (b) somewhere within the Candida albicans glucoamylase gene provided that such gene includes its signal peptide coding region, and any functional mutant or derivative thereof. The creation of the rS1d gene and its expression under the lacZ transcriptional and translational control on plasmid pTXS11 is described in full in Loch et al., *Infect. Immun.*, 55, 963-967 (1987), the entire disclosure of which is hereby incorporated by reference.

By the term 'functional mutant and derivative' is meant any protein encoded by any mutation or derivation of the S1-GA protein coding sequence which, upon expression in a host cell, e.g., a Saccharomyces host cell results in a protein which retains the biological activity of the S1-GA protein, e.g., solubility and a protective immunogenic response against pertussis. Production of such mutants and derivatives is within the routine skill of the art. Such functional mutants and derivatives of the S1-GA coding sequence include chemically modified forms thereof, e.g., glycosylated and alkylated forms which have substantially the same biological activity as S1-GA protein. They also include mutants and derivatives in which one or more amino acids have been rearranged, added, deleted or substituted or in which a

sequence of amino acids has been inserted or deleted. Thus, by the term 'functional mutants and derivatives' is meant a protein which is a mutant or derivative of the S1-GA protein coding sequence, which encodes a protein which has substantially the same biological activity as native S1-GA protein. Such functional mutants and derivatives include, but are not limited to, deletions or additions of amino acids, which encodes a protein which has substantially the same biological activity as native S1-GA protein. Such functional mutants and derivatives include, but are not limited to, deletions or additions of amino acids, which encodes a protein which has substantially the same biological activity as native S1-GA protein. Such functional mutants and derivatives include, but are not limited to, deletions or additions of amino acids, which encodes a protein which has substantially the same biological activity as native S1-GA protein.

derivatives are conventionally prepared by one of skill in the art employing, e.g., site directed mutagenesis, random mutagenesis, chemical synthesis of nucleotide or peptide sequences, truncation or deletion of part of the coding sequence for native S1-GA protein, chemical modification of the above or concatenation of more than one segment of the S1-GA protein coding sequence. Such functional mutants and derivatives of the S1-GA protein coding sequence also include chemically modified forms thereof, e.g., glycosylated and alkylated forms which have substantially the same biological activity as native S1-GA protein.

This invention is also related to a recombinant DNA molecule, such as a cDNA, which contains the S1-GA protein coding sequence. Such cDNA can be prepared by conventional techniques such as by reverse transcription of mRNA isolated by conventional techniques from a cell or culture producing S1-GA protein, such as by the method of Han et al., Biochemistry, 26, 1617-1625 (1987). As a further embodiment of the recombinant DNA molecule of the invention there is provided a recombinant DNA vector comprising the coding sequence of this invention. This vector preferably contains the above coding sequence in operative association with suitable expression control sequences. By 'suitable expression control sequences' is meant those sequences which regulate the transcription and translation of a structural gene. Such expression control sequences are well known in the art.

Still a further aspect of this invention is a host cell transformed with the vector of this invention. Such host cell is capable of growth in a culture medium and is capable of expressing the coding sequence of the invention. Such host cell is prepared by the method of this invention, i.e., transforming a desired host cell with the vector of the invention. Such transformation is accomplished by utilizing conventional transformation techniques. Host cells which may be suitable include, but are not limited to, mammalian cells insect cells, bacterial cells, plant cells and fungal cells. Thus, this invention is not limited to any specific host cells, although a Saccharomyces host cell is preferred.

The present invention also relates to a method of producing the protein encoded by the S1-GA coding sequence of this invention which comprises culturing the transformed host of the invention in appropriate culture media and isolating such protein. By "appropriate culture media" is meant that media which will enable the transformed host to survive and which will also enable such host to express the coding sequence of the invention in recoverable quantity. It will be appreciated by one of skill in the art that the appropriate culture media to use will depend upon the host cell employed. The isolation of the protein so produced is accomplished from a culture lysate of the host or, if appropriate, directly from the host's culture medium and is carried out by conventional protein isolation techniques such as those discussed above.

This invention also relates to a synthetic DNA molecule containing the S1-GA protein coding sequence. By the term 'synthetic' is meant a DNA molecule prepared by synthetic rather than recombinant techniques. Such techniques are well known in the art using commercially available DNA synthesizers.

The vaccine of this invention comprises an immunoprotective and non-toxic amount of S1-GA protein. Such vaccine preferably contains 5-25  $\mu$ g of such S1-GA protein but is not limited to these amounts.

Other antigens which are known to be desirably administered in conjunction with pertussis toxin may also be included in the vaccine of this invention. Such additional components are known to those of skill in the art. Preferable additional components include tetanus toxoid and/or diphtheria toxoid as well as filamentous haemagglutinin (FHA) and/or any other protective antigen of B. pertussis.

The provision of such a vaccine thus allows another aspect of the present invention, i.e., a method for immunizing a human against pertussis which comprises administering the vaccine of the subject invention to such human.

The mode of administration of the vaccine of the invention may be any suitable route which delivers an immunoprotective amount of the protein of the subject invention to the host. However, the vaccine is preferably administered parenterally via the intramuscular or deep subcutaneous routes. Other modes of administration may also be employed, where desired, such as oral administration or via other parenteral routes, i.e., intradermally, intranasally, or intravenously.

The vaccine of the invention may be prepared as a pharmaceutical composition containing an immunoprotective and non-toxic amount of S1-GA protein in a nontoxic and sterile pharmaceutically acceptable carrier. Where the administration of the vaccine is parenterally, the protein comprised by the vaccine can be optionally admixed or absorbed with any conventional adjuvant, for use as a non-specific irritant to attract or enhance an immune response. Such adjuvants include, among others, aluminum hydroxide, aluminum phosphate, aluminum stearate and aluminum borate, such as Quil A. As a further exemplary

embodiment of the invention, the protein comprised by the vaccine can be administered with an immunostimulating macromolecule, or a tetanus toxoid. Alternatively, an aqueous suspension or solution containing the protein of vaccine of the invention, preferably buffered at physiological pH, may be designed

for oral ingestion.

It is preferred that the vaccine of the invention, when in a pharmaceutical preparation, be present in unit dosage forms. The appropriate therapeutically effective dose can be readily determined by those of skill in the art. The effective amount of S1-GA protein contained in the vaccine of this invention may be in the range of the effective amounts of antigen in conventional *B. pertussis* acellular or component vaccines, i.e., 5-25  $\mu$ g of protein. This dose may optionally be delivered with various amounts of filamentous haemagglutinin (FHA) (about 10  $\mu$ g-25  $\mu$ g) and/or agglutinogens or other antigens. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, general health, sex, and diet of the patient; the time of administration; the route of administration; synergistic effects with any other drugs being administered; and the degree of protection being sought. Of course, the administration can be repeated at suitable intervals if necessary. Preferably, three doses will be administered, with 3 to 12 months intervals between each dose, with an optional booster dose later in time.

In the following examples, specific embodiments of the invention are more fully disclosed. Examples 1-7 illustrate the expression of *Candida* genes in *Saccharomyces* to complement the functions of the native *Saccharomyces* host cell. Examples 8 and 9 illustrate the isolation and use of *Candida* replicator regions to control expression of a *Candida* gene in *Saccharomyces*. Examples 10-13 illustrate the use of a *Candida* signal sequence as part of the expression and secretion system for other non-yeast heterologous genes in *Saccharomyces*. Examples 13 and 14 relate to the vaccine of the invention. These examples are intended to be illustrative of the subject invention and should not be construed as limiting its scope.

#### EXAMPLE 1 - ISOLATION OF DNA FROM *C. ALBICANS*

##### A. Construction of the genomic library of *C. albicans*

The DNA from *C. albicans* strain 792-1 (courtesy of J. Kwon-Chung, Natl. Inst. Health, U.S.A.) was isolated by the procedure described by Rigsby et al., cited above. The DNA purified by this procedure was on the average 100 kilobase pairs (kbp) in length.

A genomic library was constructed in the *Saccharomyces-Escherichia coli* shuttle vector, YEp13 [ATCC 37115], described by Broach et al., cited above. Calculation of the size of the genome of *C. albicans* based on the data provided by Rigsby et al., cited above, suggests that the genome size of *C. albicans* is about  $2 \times 10^4$  kbp. Therefore, some 8000 independent transformants carrying an insert of 10 kbp are required to provide a genomic library with a probability of 99% that each of the sequences of *C. albicans* will be represented at least once in this library. This was calculated by the method disclosed by Maniatis et al., "Molecular Cloning - A Laboratory Manual", Cold Spring Harbor Laboratory (1982).

The DNA was partially digested with the restriction endonuclease *Sau*3A and fractionated by centrifugation in a 5-20% sucrose gradient. The fractions containing DNA fragments in the range of 10 kbp were pooled and used for ligation. Plasmid YEp13 DNA was cleaved at the unique *Bam*HI site and then treated with alkaline phosphatase to prevent self-ligation. Equal concentrations (200 ng. in a 10  $\mu$ l reaction mixture) of *C. albicans* and YEp13 DNA were ligated.

*E. coli* strains MC1061 (Casadaban et al., J. Mol. Biol., 138:179-207 (1980)) and MB101 (Bolivar et al., Gene, 2:95-113 (1979)) were transformed with the ligated mixture according to Lederberg, and Cohen, J. Bacteriol., 119:1072-1074 (1974). With the modification that a 45 min. expression time at 37°C in Luria Broth (LB) was allowed. An estimate of the number of independent transformants was obtained by plating a small sample on LB plates containing 50  $\mu$ g/ml of ampicillin. Simultaneously, the remaining sample was inoculated into LB broth containing 50  $\mu$ g/ml of ampicillin and incubated overnight to use for DNA isolation. The plate results indicated that some 16,000 independent clones were obtained. To estimate the percentage of transformants which contained inserts, 100 transformants were tested for their resistance to ampicillin and to tetracycline. Since the *Bam*HI site is in the tetracycline resistance gene (*tet*<sup>R</sup>), transformants that contain plasmids with the inserted DNA should be sensitive to tetracycline and resistant to ampicillin, while transformants containing plasmids without inserts should be resistant to both antibiotics. The results obtained indicated that all the transformants were sensitive to tetracycline and resistant to ampicillin.

Plasmids containing inserted DNA of *C. albicans* were isolated from the transformants by the method of Maniatis et al., cited above, using an aliquot of this original culture as control.

## B. Expression of *C. albicans* genes in *S. cerevisiae*

### (1) General methodology

To isolate cloned DNA sequences of *C. albicans* by expression in *S. cerevisiae*, genetically defined haploid strains of *S. cerevisiae* were employed as hosts for DNA mediated transformation. Transformation was performed a) by the spheroplast procedure, as described by Sherman et al., "Methods in Yeast Genetics Laboratory Manual", Cold Spring Harbor Laboratory (1982), or by the lithium chloride or lithium acetate method described by Ito et al., cited above. The yeast were plated on minimal medium (Yeast nitrogen base-Difco, YNB) containing all necessary supplements except one.

Transformation of a defined *S. cerevisiae* mutant with a plasmid containing the wild type allele of the specific mutation allows the transformed host to grow under conditions restrictive to the non-transformed mutant. In the examples following, the *LEU2* gene was used as the first selectable marker. The transformed yeast were then subcultured onto additional types of media to select transformants that expressed other functions originally lacking in the *S. cerevisiae* host strain.

The plasmids were recovered from *S. cerevisiae* by making a DNA preparation of the desired transformant, using the procedure of Sherman et al., cited above, and using this DNA to transform *E. coli* by known techniques and used for further analysis. Such further study included: a) transformation of yeast to confirm the presence of a gene conferring the same phenotype as found in the original transformed yeast cell; b) hybridization analysis to confirm the presence of DNA of *C. albicans* in the hybrid plasmid; and c) further analysis of the *C. albicans* DNA fragment both structurally and functionally.

### (2) Isolation of a *Candida albicans* gene that complements the *trp1* mutation of *S. cerevisiae*.

Spheroplasts of *S. cerevisiae* strain DBY746 ( $\alpha$ , *his3-1 leu2-3, leu2-112, ura3-52, trp1-289*) were transformed with DNA isolated from the *C. albicans* library. *S. cerevisiae* strain DBY746 was obtained from Yeast Genetic Stock Center, Department of Biophysics and Medical Physics, University of California, Berkeley, California, U.S.A. The cells were plated on minimal medium (Difco YNB) containing all required supplements except leucine. The transformed yeast cells expressed the plasmid encoded *LEU2* gene. The *Leu*<sup>+</sup> cells were harvested from these plates and plated onto minimal medium lacking both leucine and tryptophan. Of the 500 transformants examined, two grew on these plates.

To test whether the growth of these two transformants on medium lacking leucine and tryptophan was due to the presence of a *C. albicans* DNA insert, rather than due to the reversion of the *trp1* mutation, the transformants were grown in non-selective yeast extract peptone dextrose (YEPD) medium for 10 generations, and then plated on minimal medium containing all supplements. The colonies were then replica plated onto minimal medium lacking only tryptophan or only leucine.

In all cases, colonies arising from cells that had lost the plasmid i.e., were *Leu*<sup>-</sup>, also required tryptophan indicating co-segregation of the *LEU2* and *TRP1* alleles.

Plasmids were recovered from the two transformed isolates by transformation of *E. coli* HB101 with small yeast DNA preparations, prepared by the method of Sherman et al., cited above, selecting for ampicillin resistant transformants. The transformed *E. coli* cells were tetracycline sensitive, indicating that the plasmid contained an insert in the *tet*<sup>R</sup> gene. The two plasmids were designated pAR84-1 and pAR84-2.

Plasmid DNA of each was prepared by the method of Maniatis et al., cited above, and used to transform *S. cerevisiae* strain DBY746. The selection of the transformed cells were performed on medium devoid of leucine but containing the supplements histidine, tryptophan and uracil. The *Leu*<sup>+</sup> transformed cells were tested for prototrophy for tryptophan. Between 88 to 100% of the transformants expressed the unselected marker *TRP*<sup>+</sup> confirming that both pAR84-1 and pAR84-2 contained a gene complementing the *S. cerevisiae* *trp1* mutation and indicating that a functional DNA sequence derived from *Candida* can be expressed in *Saccharomyces*.

To test for specific hybridization of the DNA insert in the recombinant plasmids to *Candida albicans* DNA, DNA from non-transformed and transformed *S. cerevisiae* strain DBY746 was isolated according to the procedure described by Sherman et al., cited above, for small scale preparations. The DNA of *E. coli*



with or without plasmid, was also prepared by the method of Maniatis et al., cited above.

The DNA samples were digested with the restriction enzymes HindIII and BglII. Digested DNA of C. albicans, strain 792-1, S. cerevisiae strain DBY746, plasmid YEp13, and plasmid pAR84-2 was electrophoresed on agarose gels, and transferred to a nitrocellulose filter by the procedure of Southern, J. Mol Biol., 98:503-517 (1975). The filter was then hybridized sequentially, first with <sup>32</sup>P labelled YEp13 DNA to test whether the vector used hybridized to the genomic DNA of C. albicans, and, second, with <sup>32</sup>P labelled pAR84-2 DNA to determine whether the new plasmid hybridized to the genomic DNA of C. albicans.

The nitrocellulose filters were prehybridized for 3 hrs. at 65° C with a solution containing 6 x SSC and 3 x Denhardt (1 x Denhardt is 0.02% Ficoll 400, 0.02% polyvinylpyrrolidone 360 and 0.02% bovine serum albumin) according to the method of Maniatis et al. cited above. Hybridization solution was the same except the <sup>32</sup>P labelled DNA probe prepared by nick translation according to the method of Maniatis et al., cited above, was added. Hybridization was performed for 18 hours at 65° C. After hybridization the filters were washed with 2 x SSC at 65° C. The filters were then exposed to X-ray film.

YEp13 DNA hybridized only to the genomic DNA of S. cerevisiae, whereas pAR84-2 DNA also hybridized to the genomic DNA of C. albicans. The hybridization of pAR84-2 to the S. cerevisiae genomic DNA is due to the S. cerevisiae LEU2 gene that is a part of the plasmid. The hybridization to C. albicans is due to homology of Candida insert with genomic sequences of Candida.

The size of the inserts in each plasmid was estimated by electrophoresis of restriction digests on agarose gels. pAR84-1 contains a 15 kbp fragment of C. albicans DNA, while pAR84-2 contains a 10.4 kbp fragment.

Using the same methodology as that used for the isolation of a C. albicans DNA fragment that complements the S. cerevisiae trp1 mutation, plasmids carrying other functional C. albicans genes have been isolated and characterized as described in the following Examples 2-7.

## EXAMPLE 2 - COMPLEMENTATION OF THE S. CEREVISIAE his3 GENE WITH C. ALBICANS DNA

The his3 of S. cerevisiae encodes imidazoleglycerol phosphate dehydrogenase. The isolation of a C. albicans gene that complements the his3 mutation of S. cerevisiae was carried out according to the procedure of Example 1. The Leu<sup>+</sup> yeast transformants were plated on medium lacking leucine and histidine. Colonies which arose were selected.

Plasmids containing a C. albicans gene that complements the his3 mutation of S. cerevisiae were recovered from the selected colonies, described above, by transformation of E. coli HB101 according to the method of Example 1. Plasmid DNA was isolated from the E. coli transformants according to the method of Example 1, and was designated pAR84-3.

Plasmid pAR84-3, containing the C. albicans gene that complements the his3 mutation of S. cerevisiae was transformed into S. cerevisiae strain DBY746 according to the procedure of Example 1.

Host S. cerevisiae microorganisms were transformed with plasmid pAR84-3 and selected according to the methods described above. These microorganisms were then tested for expression of the his3 gene.

## EXAMPLE 3 - EXPRESSION OF THE C. ALBICANS BENOMYL RESISTANCE GENE IN S. CEREVISIAE

The C. albicans benomyl resistance gene confers high level resistance to the anti-fungal agent benomyl (methyl 1-(butylcarbamoyl)-2-benzimidazole-carbamate). Strains of C. albicans are more resistant to benomyl than S. cerevisiae. This gene thus confers a new characteristic to a transformed S. cerevisiae strain, as follows.

The isolation of the C. albicans benomyl resistance gene was carried out according to the procedure of Example 1, except that S. cerevisiae strain JWG2-2C ( $\alpha$ , leu2-3, leu2-112) was employed as the host. S. cerevisiae strain JWG2-2C was derived from a cross of S. cerevisiae strain DC5 (leu2-3, leu2-112, his3, gal2, can1-11), with S. cerevisiae strain YM146 ( $\alpha$ , gal<sup>+</sup>  $\Delta$ 152, trp1-289, ura3-52, rho<sup>+</sup>). S. cerevisiae strain DC5 was obtained from James R. Broach, S.U.N.Y., Stonybrook, New York, U.S.A. S. cerevisiae strain YM146 was obtained from Mark Johnson, Washington State University, Pullman, Washington, U.S.A. The

transformants were tested for benomyl resistance. Plasmid DNA was isolated from individual E. coli transformants according to the small scale method of Sherman et al., cited above. The size of the C.

albicans insert in the recovered plasmids was estimated by agarose gel electrophoresis.

The plasmid DNA containing the *C. albicans* benomyl resistance gene was isolated from the *E. coli* transformants, and used to transform *S. cerevisiae* strain JWG2-11D ( $\alpha$ , his3, trp1-289, leu2-3, leu2-112, gal $\Delta$ 152) according to the procedure of Example 1. *S. cerevisiae* strain JWG2-11D was derived by the same method as *S. cerevisiae* strain JWG2-2C.

Host *S. cerevisiae* microorganisms, transformed with a plasmid containing the *C. albicans* benomyl resistance gene according to the method described above, were selected as described above. The presence of plasmid in the transformed hosts was confirmed according to the method of Example 1 by plating on medium containing benomyl. Three independent isolates, designated pCABen17, pCABen3 and pCABen15, were further analyzed by restriction mapping, and all were found to contain an identical 5.8 kilobase pair (kbp) DNA fragment from *C. albicans*.

#### EXAMPLE 4 - EXPRESSION OF THE *C. ALBICANS* SORBITOL UTILIZATION GENE IN *S. CEREVISIAE*

The *C. albicans* sorbitol utilization gene allows for growth on the carbon source sorbitol. Strains of *C. albicans* can grow on sorbitol but *S. cerevisiae* and other species of *Saccharomyces* cannot utilize this sugar. This gene thus confers a new characteristic to the transformed *S. cerevisiae* strain, as follows.

The isolation of the *C. albicans* sorbitol utilization gene was carried out according to the procedure of Example 1, except that *S. cerevisiae* strain JWG2-2C was employed as the host. Leu<sup>+</sup> transformants were plated on minimal medium (Difco YNB) containing all required supplements except leucine and glucose, and also containing 2% sorbitol.

Plasmids containing a *C. albicans* gene that codes for an enzyme conferring sorbitol utilization ability were recovered according to the procedure for Example 1. Plasmid DNA was isolated from individual *E. coli* transformants according to the small scale method of Sherman et al., cited above, and the size of the *C. albicans* insert of the recovered plasmids was estimated by agarose gel electrophoresis.

Plasmid DNA containing the *C. albicans* sorbitol utilization gene was isolated from the *E. coli* transformants described above, and used to transform *S. cerevisiae* strain JWG2-11D according to the method of Example 1.

Host cells of *S. cerevisiae* transformed with a plasmid containing the *C. albicans* sorbitol utilization gene were selected as described above. The presence of the plasmid in the transformed hosts was confirmed according to the method of Example 1. DNA of two independent isolates, designated pCAsor2 and pCAsor9, were further analyzed by restriction mapping. pCAsor2 was found to contain a 4.5 kbp DNA insert from *C. albicans*, while pCAsor9 was found to contain a 4.8 kbp DNA insert from *C. albicans* which showed little homology to the pCAsor2 *C. albicans* DNA insert.

#### EXAMPLE 5 - EXPRESSION OF THE *C. ALBICANS* GALACTOKINASE GENE IN *S. CEREVISIAE*

The isolation of the *C. albicans* galactokinase gene, which complements a *S. cerevisiae* gal mutation, was carried out according to the procedure of Example 1, except that *S. cerevisiae* strain JWG2-11D was employed as the host. Leu<sup>+</sup> transformants were plated on minimal media (Difco YNB) containing all required supplements except leucine and glucose, and also containing 2% galactose.

Plasmids containing the *C. albicans* gene that codes for galactokinase were recovered in *E. coli* according to the procedure for Example 1. Plasmid DNA was isolated from individual *E. coli* transformants according to the method of Example 1.

The plasmid DNA containing the *C. albicans* galactokinase gene was isolated from the *E. coli* transformants as described above and used to transform *S. cerevisiae* strain JWG2-11D according to the method of Example 1.

Host *S. cerevisiae* microorganisms, transformed with a plasmid containing the *C. albicans* galactokinase gene were selected as described above, and the presence of the plasmid in the transformed host was confirmed according to the method of Example 1. Two independent isolates, designated as pCAGal1 and pCAGal2, were further analyzed by restriction mapping, which revealed that the 7 kbp *C. albicans* DNA

insert was identical in both isolates. The identity of the insert as the *C. albicans* galactokinase gene was further demonstrated by measuring the level of galactokinase activity in strains JWG2-11D transformed with plasmid pCAGal1-4. Enzyme activity measured by the method of Butt et al. *PNAS-USA* 71: 3332-3336 (1984), was only detected in galactose induced cultures containing pCAGal1-4.

The promoter of the *C. albicans* galactokinase gene was demonstrated to be positively activated by the *S. cerevisiae* positive regulator, the GAL-4 gene product, as follows. Strain JWG2-11D containing pCAGall-4 was mated to another *S. cerevisiae* strain (genotype:  $\alpha$ -trp1-289, leu2-3, leu2-112, gal1-152, ura3-52) containing a 2  $\mu$ m Leu2 shuttle vector containing the *S. cerevisiae* GAL-4 gene. The resulting diploid was maintained on medium lacking both leucine and tryptophan. The induced level of galactokinase was significantly higher in the presence of the plasmid expressing the GAL-4 gene. This result indicates that the *C. albicans* sequence in pCAGall-4 contains a galactose regulated promoter which is responsive to *S. cerevisiae* regulatory proteins.

#### EXAMPLE 6 - EXPRESSION OF THE *C. ALBICANS* ISOMALTASE GENE IN *S. CEREVISIAE*

The enzyme isomaltase cleaves  $\alpha$ -1-6-glucosidic bonds, and enables microorganisms expressing it to grow on the carbon source known as  $\alpha$ -methyl-D-glucopyranoside (e.g. isomaltose). Strains of *C. albicans* can grow on this substrate but most strains of *S. cerevisiae* cannot naturally utilize this sugar. This enzyme has been found in other species of *Saccharomyces*. Thus, the *C. albicans* gene which codes for isomaltase confers a new characteristic to the transformed *S. cerevisiae* host.

The isolation of the *C. albicans* isomaltase gene was carried out according to the procedure of Example 1. *Leu*<sup>+</sup> transformants were plated on minimal media (Difco YNB) containing all required supplements except leucine, and also containing 2%  $\alpha$ -methyl-D-glucopyranoside.

Plasmids containing the *C. albicans* gene that codes for isomaltase were recovered according to the procedure of Example 1. Plasmid DNA was isolated from individual *E. coli* transformants according to the method of Example 1.

The plasmid DNA containing the *C. albicans* isomaltase gene was isolated from the *E. coli* transformants described above and used to transform *S. cerevisiae* strain JWG2-11D according to the method of Example 1.

Host *S. cerevisiae* microorganisms, transformed with a plasmid containing the *C. albicans* isomaltase gene were selected as described above, and the presence of the plasmid in the transformed host was confirmed according to the method of Example 1.

#### EXAMPLE 7 - EXPRESSION OF THE *C. ALBICANS* GLUCOAMYLASE GENE IN *S. CEREVISIAE*

The *C. albicans* glucoamylase gene codes for an enzyme allowing for growth on soluble starch. Strains of *C. albicans* can utilize such soluble starch as a substrate, but *S. cerevisiae* cannot naturally utilize this carbon source. This enzyme, glucoamylase, has been found in other species of *Saccharomyces*. Thus, the *C. albicans* gene which codes for glucoamylase (which cleaves  $\alpha$ -1-4 glucosidic bonds) confers a new characteristic to the transformed *S. cerevisiae* host.

The isolation of the *C. albicans* glucoamylase gene was carried out substantially according to the procedure of Example 1, except that *S. cerevisiae* strain JWG2-2C was employed as the host. *Leu*<sup>+</sup> transformants were plated on minimal media (Difco YNB) containing all required supplements except leucine and glucose, and also containing 2% soluble starch. The *E. coli* transformants were probably secreting the glucoamylase since satellite colonies began to grow adjacent to the original large colonies.

Plasmids containing the *C. albicans* glucoamylase gene were recovered in *E. coli* and plasmid DNA was isolated from individual *E. coli* transformants according to the methods of Example 1.

Plasmid DNA containing the *C. albicans* glucoamylase gene isolated from the *E. coli* transformants as described above, was used to transform *S. cerevisiae* strain JWG2-11D according to the method of Example 1.

Host *S. cerevisiae* microorganisms, transformed with a plasmid according to the method described above, were selected as described above. The presence of the plasmid in the transformed host was confirmed according to the method of Example 1. Three independent isolates, designated as pSTA27-1, pSTA1-3 and pSTA15-1, were further analyzed by restriction mapping which revealed that pSTA27-1 and pSTA1-3 contain an identical *C. albicans* DNA insert while pSTA15-1 contains a possibly different *C.*

plasmid. The *S. cerevisiae* strains were also transformed with the parent vector plasmid strain (genotype  $\alpha$ , leu2, his1). The same three strains were also transformed with the parent vector plasmid YEp13 lacking any *C. albicans* sequences. The growth of all six transformed strains was tested on complete

(YEP) medium containing 0.5% maltooligosaccharides (Mos) (Pfanstiel - Laboratories Inc.) as the sole carbon source.

This compound, Mos, consists of a mixture of glucose polymers linked by alpha-1,4-glucosidic bonds. It is devoid of glucose, maltose and maltotriose, and thus consists of higher order oligomers which cannot be used as carbon sources by *Saccharomyces cerevisiae* strains. This is due to the fact that: 1) these polysaccharides do not enter *S. cerevisiae* cells and 2) *S. cerevisiae* strains do not secrete an enzymatic activity (amylase or glucoamylase) capable of degrading Mos extracellularly.

While all three strains transformed with plasmid YEp13 failed to grow on Mos containing medium, the same three strains carrying plasmid pSTA27-1, as well as *C. albicans* strain 792-1 (courtesy of J. Kwon-Chung, Natl. Inst. Health, U.S.A.) grow well on this medium. This result indicates that the *C. albicans* DNA fragment cloned in pSTA27-1 carries a gene encoding an enzyme capable of degrading Mos into a usable carbon source.

This gene is expressed in *S. cerevisiae*, and its product is secreted into the culture medium.

In order to confirm the expression and secretion of the *C. albicans* glucoamylase by a *S. cerevisiae* host transformed therewith, the culture media of all six transformed *S. cerevisiae* strains, as well as that of *C. albicans* strain 792-1, were tested for the presence of glucoamylase activity according to the method described by Searle et al., *FEMS Microbiology, Letters*, 111, 211-212 (1981). It was found that while the culture media of *S. cerevisiae* strains transformed with YEp13 was devoid of any detectable glucoamylase activity, the culture media of *C. albicans* strain 792-1 and that of *S. cerevisiae* containing pSTA27-1 contained appreciable amounts of glucoamylase. In fact, it was found that *S. cerevisiae* strains carrying plasmid pSTA27-1 secrete 15 to 30 fold more glucoamylase activity into their culture medium than does *C. albicans* strain 792-1.

The glucoamylase activity secreted by *S. cerevisiae* strains transformed with pSTA27-1 was further characterized and compared to that secreted by *C. albicans* strain 792-1 in the following manner:

a. The pH optimum of the enzyme secreted by *S. cerevisiae* strains was determined to be between 4.5 and 6.0 and is identical to that of the enzyme secreted by *C. albicans*.

b. The temperature optimum of the *S. cerevisiae* secreted enzyme was found to be 60°C and is identical to that of the *C. albicans* secreted enzyme.

c. Aliquots of the culture medium derived from an *S. cerevisiae* strain (genotype:  $\alpha$ , *leu2*, *his1*), described above, transformed with either YEp13 or pSTA27-1 and of *C. albicans* strain 792-1 were analyzed by SDS polyacrylamide gel electrophoresis. A diffuse protein band of a molecular weight higher than 94K dalton is present in the above-described *S. cerevisiae* strain transformed with pSTA27-1 medium but absent from medium derived from that strain transformed with YEp13. A similar, if not identical, protein band is detected, although in much lower amounts, in medium derived from *C. albicans* strain 792-1.

Taken together, these results indicate the fragment of *C. albicans* DNA cloned in pSTA27-1 carries a *C. albicans* gene encoding a glucoamylase that this gene is expressed in the yeast *S. cerevisiae*, and that the product of this gene is secreted into the culture medium by *S. cerevisiae* cells.

Examples 8 and 9 report the isolation and use of several *C. albicans* replicator regions that support autonomous replication of recombinant plasmids in heterologous host *S. cerevisiae* microorganisms.

#### EXAMPLE 8 - TRANSFORMATION OF *S. CEREVISIAE* WITH A RECOMBINANT PLASMID CONTAINING A *C. ALBICANS* REPLICATOR REGION

##### A. Materials and methods

*Escherichia coli* strain SF<sub>8</sub> (*C600*, *leuB*, *thr*, *pro*, *Bl*, *rk*<sup>-</sup>, *mk*<sup>-</sup>, *recBC*<sup>-</sup>, *lopII*, *lig*<sup>+</sup>) [Cameron et al., *PNAS, USA*, 72, 3416-3420 (1975)], was used for bacterial transformation and propagation of plasmids. Bacteria were grown on LB (rich media) or M9 (minimal media) supplemented as required [(Miller, "Experiments In Molecular Genetics", Cold Spring Harbor Laboratory, New York, 431-433 (1972)). *Saccharomyces cerevisiae* strain 483 (genotype:  $\alpha$ , *leu2-3*, *leu2-112*, *his4-519*, *can1a*) was used as host for

Plasmid M851 [Gorman et al., *Mol. Gen. Genet.*, 183, 306-313 (1981)] was employed. *C. albicans* strain 792-1 was used as the source of genomic DNA.

Covalently closed circular plasmid DNA was prepared from M9-glucose grown *E. coli* cultures amplified

with chloramphenicol. DNA was extracted and purified by a cleared lysate technique essentially as described by Clewell, J. *Bacteriol.*, 110, 667-676 (1972).

Crude *E. coli* plasmid preparations, used for yeast transformation and preliminary gel analysis, were made using the procedure of Birnboim and Doly, *Nucl. Acids Res.*, 7, 1513-1523 (1979). Ten ml of bacterial culture ( $2 \times 10^{10}$  cells) yielded 1-2  $\mu$ g of crude plasmid DNA. For rapid screening of bacterial transformants for the presence of plasmid, single colony mini-lysates, prepared by the same method, were used.

For recovery of total DNA from *S. cerevisiae*, the small scale method of Sherman et al., cited above, was employed.

*C. albicans* DNA was prepared by the method of Rigby et al., *J. Mol. Biol.*, 113, 237-251 (1977).

All gel analysis and hybridization procedures employed were according to the method of Maniatis et al., cited above. *E. coli* were transformed using the calcium-heat shock method, as modified by Dagert and Ehrlich, *Gene*, 6, 23-28 (1979). The procedure used for yeast transformation was essentially as described by Beggs *Nature*, 275, 104-109 (1978), using  $\beta$ -glucuronidase-sulfatase (Sigma) for spheroplast formation, with the exception that a lower concentration of plasmid DNA (0.5-1.0  $\mu$ g per  $10^8$  spheroplasts) was used.

## B. Results

### 1. Plasmid Construction

Total *Candida* DNA, digested to completion with restriction endonuclease *Pst*I, was mixed with *Pst*I-cut M851 plasmid DNA at concentrations estimated to give maximal recovery of plasmids containing *Candida* inserts by the method of Dugaiczky et al., *J. Mol. Biol.*, 96, 171-184 (1975), and ligated with T4 polynucleotide ligase. This procedure inserted the DNA by ligation into the unique *Pst*I site of M851. The ligated mixture was used to transform *E. coli* strain SF8 to leucine prototrophy. *Leu*<sup>+</sup> transformants were then screened for ampicillin sensitivity. Interruption of the *amp*<sup>r</sup> gene by an inserted fragment results in loss of drug resistance. Thus *amp*<sup>s</sup> colonies (5.8% of the *Leu*<sup>+</sup> bacterial transformants) were selected as presumably harboring plasmids containing sequences from *Candida*. These hybrid plasmids were designated pCA followed by an isolate number.

### 2. Transformation of Yeast

Plasmids containing a yeast replicator sequence transform yeast at high frequency, reflecting autonomous replication of the plasmid in the eucaryotic cell. If the plasmid lacks a functional replicator it also will transform, but at a low frequency. Presumably, this low-frequency stable transformation reflects integration of the plasmid *leu2*<sup>+</sup> gene into chromosomal DNA by homologous recombination. If *Candida* chromosomal replicators are functional in yeast, hybrid plasmids containing them might be characterized by the ability to transform yeast at high frequency.

Forty-one (41) pCA plasmids were individually tested for their ability to transform *leu2 S. cerevisiae* to *Leu*<sup>+</sup>. Five of the forty-one pCA plasmids tested showed high frequency transformation. These results indicate that such plasmids can replicate autonomously in *S. cerevisiae*.

### 3. Detection of Plasmid DNA in *S. cerevisiae*

If high frequency transformation indicates the presence of a *Candida* sequence that allows autonomous plasmid replication, it should be possible to recover covalently closed circular molecules from the unstable *Leu*<sup>+</sup> yeast transformed with DNA from the pCA plasmids.

Total yeast DNA was extracted from parental and transformed yeast strains, electrophoresed in a 1.0% agarose gel, and transferred to nitrocellulose paper. The DNA was then hybridized with P<sup>32</sup>-labelled pBR322 probe. This probe has no homology with yeast chromosomal DNA. The DNA from transformed cultures was

#### 4. Transformation of *E. Coli* with Yeast Plasmid DNA

The yeast plasmid preparations analyzed by hybridization techniques were also examined for ability to transform *leuB*, *E. coli* to *Leu*<sup>+</sup>. Bacterial colonies selected on leucine deficient medium were screened for the presence of plasmid by agarose gel electrophoresis of mini-lysates. The five yeast extracts that were shown to contain plasmids by hybridization have rise to *Leu*<sup>+</sup> *E. coli* that contained plasmid DNA. Plasmid DNA of one of the five high frequency transformation pCA plasmids was compared with the plasmid isolated from *E. coli* following transfer through the *S. cerevisiae* host. The plasmids, containing a 4.0 kilobase pair (kbp) *C. albicans* insert, were found to be identical.

Other replicator regions of *Candida* useful in the autonomous maintenance of plasmids used to transform *Saccharomyces* can be isolated according to the method of Kawamura et al., cited above; Hsu et al., cited above; or Tikhomirova et al., cited above. Plasmids containing such replicator regions can be prepared according to the method described in this Example.

#### EXAMPLE 9 - TRANSFORMATION OF *S. CEREVISIAE* WITH A PLASMID CONTAINING A *C. ALBICANS* REPLICATOR REGION AND A *C. ALBICANS* CODING SEQUENCE

A plasmid containing a *C. albicans* replicator region and a *C. albicans* coding sequence can be prepared by combining the methods of Examples 1 and 8. Such a plasmid can be used to transform *S. cerevisiae* according to the method of Example 1, and expression of the *C. albicans* coding sequence by the transformed *S. cerevisiae* host can be detected according to the method of Example 1.

Genes for glucoamylases from mouse (Thomsen, 1983, *Carlsberg Res. Commun.*, 48:545), wheat, *Asperigillus* (Innis et al., 1985, *Science*, 228:21), and *S. diastaticus* (Yamashita and Fukui, 1983, *Agric. Biol. Chem.*, 47:2689) have been shown to be secreted into the medium when expressed in *S. cerevisiae*. When produced by *C. albicans* or by a transformed *S. cerevisiae* host, the *C. albicans* enzyme glucoamylase is excreted into the medium. It has now been found that the *C. albicans* glucoamylase gene, also called the STA gene, isolated as described in Example 7 provides DNA sequences for a novel promoter and a signal peptide which functions in protein secretion. The use of a signal peptide for secretion of a cytoplasmic protein is disclosed, for example, see Silhavy, et al., U.S. Patent 4,335,336 issued June 22, 1982. This glucoamylase gene signal peptide can be used in a system for secretion of native or heterologous proteins expressed by *Candida*, *Saccharomyces* or other yeast cells transformed therewith. This expression system targets the heterologous protein to the secretion pathway leading to post- or co-translational modifications of the synthesized protein.

Both the promoter and signal sequences of *C. albicans* glucoamylase gene are identified in Fig. 1. Fig. 3 illustrates the signal peptide alone. The signal sequence enables the targeting of the heterologous protein to the secretion pathway. Thus, it may be used in an expression system to enable secretion of a recombinant protein into the culture medium whether under control of its own promoter or under control of heterologous yeast promoters.

The following Examples 10-13 demonstrate a variety of uses of this functional DNA sequences of *Candida albicans*, e.g. the use of the *C. albicans* promoter and signal sequence to express and secrete a heterologous protein in *S. cerevisiae* and the use of the *C. albicans* signal sequence with other yeast promoters for such expression and secretion.

In general in these examples *E. coli* vectors are constructed which allow inframe translational fusion of heterologous proteins or peptides to the promoter/signal peptide region of this glucoamylase gene. These hybrid gene constructions, (e.g., glucoamylase promoter and signal peptide regions or the signal peptide under control of another yeast promoter) fused to a heterologous gene are then transferred preferably into *E. coli*/*Saccharomyces* shuttle plasmid vectors for transformation into yeast cells.

#### EXAMPLE 10 - EXPRESSION AND SECRETION OF HIV-I GP160 UNDER CONTROL OF *C. ALBICANS* GLUCOAMYLASE PROMOTER/SIGNAL PEPTIDE REGION IN *S. CEREVISIAE*

Interest for use in potential AIDS vaccines and therapeutics has been expressed in the use of HIV-I proteins for detection of HIV infection.

#### A. Construction of *E. coli* plasmid vectors

*E. coli* plasmid pUC13 [J. Viera et al., *Gene*, 19:259-68 (1982)] was linearized by co-digestion with restriction enzymes *AccI* and *HindIII*. The linearized plasmid was ligated to a *Tag1* to *HindIII* DNA fragment constructed as follows: A *Tag1* to *Fnu4H1* DNA fragment comprising the promoter and signal peptide region of the yeast *Candida albicans* *STA* gene (encoding the secreted enzyme, glucoamylase) was purified from the cloned gene. The nucleotide sequence of this DNA fragment is shown in Fig. 1.

A synthetic double stranded oligonucleotide having the following sequence:

***Fnu4H1***

***BglII***

***HindIII***

**CGCTCCAGATCTA**

**CGAGGTCTAGATTCGA**

was ligated to the *Tag1* to *Fnu4H1* fragment described above. The product of this ligation, (a *Tag1* to *HindIII* DNA fragment) after gel purification, was itself ligated to the *AccI*/*HindIII* digested pUC13 DNA. The ligation mixes were used to transform *E. coli* strain MM294 (available from the American Type Culture Collection as ATCC 33625 or ATCC 39607).

Transformants were selected on LB plus ampicillin plates and analyzed by colony filter hybridization using a radiolabelled probe containing sequences homologous to the *Tag1* to *Fnu4H1* fragment. Plasmid DNA from appropriate transformants was subjected to restriction enzyme and nucleotide sequencing analyses. The selected plasmid resulting from these analyses pTI-18 (pRIT13046) contains the entire promoter region of the yeast *STA* gene, as well as the DNA sequence coding for the first 22 NH<sub>2</sub> terminal amino acids of the glucoamylase protein, which represent the signal peptide of this secreted protein. Immediately following the signal peptide cleavage site, a unique *BglII* restriction cleavage site has been introduced as part of the synthetic oligonucleotide described above.

Cloning of the coding sequence (or part thereof) of foreign genes into the *BglII* site in such a way as to respect the reading frame of both the signal peptide as well as that of the foreign protein permits once the construct is transferred into a yeast plasmid and introduced into yeast cells, the expression of the hybrid gene via the glucoamylase *STA* gene promoter and the targeting of the foreign protein to the secretion pathway via the glucoamylase signal sequence.

#### B. Cloning of the HIV envelope glycoprotein (gp160) coding sequence in a yeast vector

A genomic clone of the HIV variant BH10 [L. Ratner et al., *Nature*, 313:277-284 (1985)] was digested with restriction enzymes *Asp718* and *XbaI*. An *Asp718* to *XbaI* DNA fragment of approximately 2550 base pairs was gel purified. This fragment encompasses the coding sequence of the envelope glycoprotein gp160 (the *env* gene), starting at amino acid Val 42 and extends several hundred nucleotides beyond the translational stop codon (TAA) of the *env* gene. This fragment was treated with Mung bean nuclease according to Maniatis et al., cited above to render it blunt ended.

Plasmid pTI-18 (pRIT13046) described above was linearized by digestion with restriction enzyme *BglII*, treated with Klenox polymerase I in the presence of deoxynucleotide triphosphates (dNTP's) in order to fill in protruding 5' ends. The fragment and plasmid were ligated using T4 DNA ligase. *E. coli* cells were transformed with the ligation mix. Transformants were analyzed by colony filter hybridization. Restriction enzymes and DNA sequence analysis was performed on plasmid DNA extracted from selected transformants.

These analyses allowed the selection of transformants containing plasmid pTI-18/gp160 (pRIT13047). This plasmid contains the HIV *env* gene fragment defined above, fused in frame to the *STA* gene signal sequence and preceded by the *STA* gene promoter region. The DNA and amino acid sequence of the relevant junction between the signal sequence coding region and the *env* gene coding sequence is provided below.

5

10

20

25

### C. Expression of the HIV env gene in yeast and localization of the gene product (gp160)

35

41

5



material by immunoblotting confirms this hypothesis.

#### D. Detection of HIV gp160 in a crude membrane extract

Fraction S2-30 was analyzed by capture ELISA for the presence of HIV-env antigenic sites. A sheep antibody to a synthetic peptide of HIV gp41 (Biochrom #D7323) was absorbed to NUNC Immunoplate 1 microtiter plates at a concentration of 5 µg/ml in PBS; 50 µl/ml of antibody was incubated overnight at 4 °C. The plates were washed by 6 fold replacement of the content of each well with PBS and 0.1% Tween 20 and incubated with 100 µl of saturation buffer (PBS + 0.1% Tween 20 + 1% bovine serum albumin + 4% newborn calf serum) for 1 hour at 37 °C. The plates were then washed as above and the individual wells were incubated with serial dilutions of S2-30 in PBS + 1% Triton. Fifty µl of antigen were incubated for 2 hours at room temperature. The plates were washed as above and incubated with a monoclonal antibody to HIV gp120 (Dupont #9284/7054). A concentration of 1 µl/ml of antibody in saturation buffer was used. Fifty µl well were incubated for 1 hr at 37 °C. The plates were washed as described and the binding of the monoclonal anti-HIV gp120 antibody was revealed using biotin labeled anti-mouse immunoglobulin reagent (Amersham RPN 1021) and streptavidin-peroxydase complex (Amersham RPN 1051), according to the manufacturer's instructions. The titration curves obtained indicated the presence of HIV env gene antigenic sites.

#### E. Optimization of HIV gp160 yield

The protein gp 160 was produced by batch fermentation of the *S. cerevisiae* strain (genotype: Mata ura3 leu2 adel gln1) containing plasmid pQU63/gp160 (pRIT13049) lasting 25-30 hours in a standard stirred tank vessel maintained at 30 °C. The rich medium (1% yeast extracted 2% peptone, 2% galactose - YEP) was maintained at pH 5.0 by base addition. The inoculum (10% v/v) was grown up for 24 hours on YEP and 2% glucose at 30 °C.

Yeast growth was rapid and was measured by reference to its Optical Density (O.D.<sub>650</sub>) at 650nm. The maximum doubling time during exponential phase was approximately 3 hours.

During the exponential growth phase, gp160 was synthesized rapidly and was measured after approximately 12 hours of growth using capture ELISA as described above. Synthesis reached a maximum of 800 µg/ml of fermentation broth at approximately 20 hours, whereupon a rapid decline in gp160 was detectable. The onset of the decline gp160 concentration coincided with the depletion of substrate galactose.

Galactose metabolism was followed indirectly by mass spectrometric measurement (each 20 seconds) of the CO<sub>2</sub> and oxygen concentration in the fermentor off-gas. The exhaustion of galactose was correlated with the rapid decline of % CO<sub>2</sub> in the fermentor off gas at approximately 20 hours.

Thus, harvesting the culture shortly before the decline of % CO<sub>2</sub> in the fermentor may be crucial for obtaining a good yield of gp160 in *S. cerevisiae*.

#### F. Purification of gp160 and micelle preparation

One volume of transformed yeast cells were ground in a Dynomil apparatus in the presence of one volume of buffer consisting of 25mM phosphate pH 7.0 containing 150mM NaCl, 1mM ethylenediaminetetraacetic acid (EDTA) and 1mM PMSF (Buffer A). The suspension is centrifuged for 10 min. at 10000xg and the pellet consisting of unbroken cells and cells debris is discarded. The supernatant is in turn centrifuged 30 min. at 30000xg and the pellet is extracted with 1% Triton X-100 in the original volume of Buffer A. After centrifugation at 30000xg (30 min) the gp160 containing supernatant is treated with 20% Polyethyleneglycol-400 to precipitate gp160. The PEG pellet is dissolved in 50mM Tris buffer pH 7.2 containing 150mM NaCl and 1% Triton X-100 and supplemented with 1.5 M CsCl and 4M guanidine

hydrochloride. The sample is ultracentrifuged in a Titanium 50.2 Rotor during 65 hours at 45000 rpm. The supernatant is then layered on a 10-30% sucrose gradient and centrifuged for 18 hours at 45000 rpm.

Fractions that become Elisa positive after Triton treatment are pooled and used for the subsequent purification of HIV-gp160 envelope glycoprotein obtained via the

This yeast synthesized HIV-gp160 envelope glycoprotein obtained via the expression/secretion system of the present invention has potential as a vaccine candidate. The post-

translational modifications allowed by this expression system effect the tertiary structure of the protein, which may, in turn, effect the induction of a relevant immune response, since a significant part of the antibody is directed to conformational epitopes. The high cysteine content and the frequent occurrence of potential N-linked glycosylation sites in gp160 suggests the formation of disulfide bridges and the further  
 5 modification of antigenic properties of the molecule by glycosylation. Thus targeting of gp 160 to the secretion pathway in an expression system using the glucoamylase signal peptide yields a molecule that appears to be structurally and antigenically more related to the native envelope glycoprotein spike. Those structural features of the env-gp160 glycoprotein obtained via the expression system of the present invention are lacking in the envelope protein expressed via non-secretion systems in yeast [see, e.g., P.J.  
 10 Barr et al., cited above] or in E. coli [see, e.g., S.D Putney et al., Science, 234:1392-96 (1986)].

For example, in this expression system, the yeast synthesized gp160 remains unprocessed into its gp120 and gp41 glycoprotein components. Thus, both proteins remain membrane associated. This is unlike the situation reported in mammalian cells [see, e.g., M.P. Kieny et al., Biotechnology, 4:790-795 (1986)] where the cleavage between gp41 and gp120 results in dissociation of the two molecules and release of  
 15 most gp120 molecules in the culture supernatant. The association to cell membranes via a hydrophobic transmembrane region of gp41 can serve for the incorporation into particulate structures such as micelles, into liposomes by association to lipids or into immunostimulating complexes (ISCOMS) by association to the detergent quill A. This can increase the immunogenicity of the molecules, both by the adjuvant properties of liposomes and ISCOMS and by allowing an oriented presentation at the surface of these  
 20 structures, similar to the viral surface.

#### EXAMPLE 11 - EXPRESSION AND SECRETION OF TGF- $\alpha$

25

##### A. Construction of basic yeast expression vectors

The plasmids used to construct vectors for TGF- $\alpha$  secretion using the glucoamylase signal sequence are based on the high copy yeast shuttle vector pYSK102 [Rosenberg et al., Genetic Engineering, 8:151  
 30 (1986)]. All contain an 823 bp EcoRI-PstI fragment of the yeast trp1 gene [Tsumper and Carbon, Gene, 10:157 (1980)], which provides a selectable marker, a 2.0 kb EcoRI-PstI fragment of the yeast 2-micron circle, [Hartley and Donaldson, Nature, 286:860 (1980)] which provides an origin of replication in yeast, a portion of pBR322 [in "Cloning Vectors", Pouwels, Enger, Valk, Eds., (1985)] which provides for replication and selection in E. coli and a yeast promoter-terminator module inserted between the HindIII and BamHI  
 35 sites of pBR322. The pBR322 sequences between BamHI and PvuII have been deleted.

The promoter employed in these constructs is either a 431 bp BamHI-RsaI fragment of CUP1 [Karin et al., PNAS USA, 81:337 (1984)], derived as a BamHI-EcoRI fragment from pYSK12 [Butt et al., Proc. Natl. Acad. Sci. USA, 81:3332 (1984); Rosenberg et al., *supra*; Gorman et al., Gene, 48:13 (1986)] or a 1.1 kb  
 40 TDH3 promoter sequence [Holland and Holland, J. Biol. Chem., 255:2596 (1980)], derived as a BamHI-EcoRI fragment from plasmid pYSK18 [Rosenberg et al., *supra*; Gorman et al., *supra*]. The transcriptional terminator is a 361 bp KpnI-HindIII of the (CYC1 gene [Smith et al., Cell, 16:753 (1979)].

A synthetic linker, with the sequence 5' GAATTCT-CGAGCCATGGCCAGTCGACGTAGTTAAGTAGGTACC 3' is present in the region between the promoter and the CYC1 termination fragment. This synthetic sequence contains the unique restriction sites XhoI, NcoI  
 45 and Sall upstream of translational stop codons in all three reading frames.

##### B. Construction of pYSK140 and pYSK142

50 The site of signal cleavage of the glucoamylase gene was determined by amino terminal amino acid sequencing of the secreted glucoamylase of Example 7. The DNA sequence of the 5' end of the glucoamylase gene coding region and the corresponding amino acid sequence, with the site of processing,

signal sequence ends after the last asterisk. This double standard fragment was inserted in the basic vectors described above which had been cleaved with XhoI and NcoI to give pYSK140 (CUP1 promoter) and pYSK142 (TDH3 promoter).

The current synthetic glucoamylase signal sequence contains the codons for the amino acids Asn Ala Ala Pro at the 3' end to provide the processing site (see Fig. 3). The presence of the two amino acids Ala and Pro at the 3' side of the cleavage site may be essential for proper cleavage. However, this results in the presence of two extra amino acids on the 5' end of any protein secreted by utilization of this signal sequence. Plasmids are being constructed which will allow fusions directly after the Asn Ala codons. Secretion and processing at the 5' end of fusion proteins made with these vectors are expected to be as useful as the present signal sequence.

#### 10 C. Construction of pYSK151 and pYSK152

A synthetic oligonucleotide corresponding to the CDNA sequence of the mature human transforming growth factor (TGF- $\alpha$ ) [Derynck et al., Cell, 38:287 (1984)] plus additional bases at the 5' and 3' end was synthesized and four conservative base changes made to provide restriction endonuclease sites. This sequence is shown in Fig. 5. The NcoI to Sall portion of the TGF- $\alpha$  sequence was inserted into NcoI and Sall cleaved pYSK140 and pYSK142 to give pYSK151 and pYSK152 respectively. The resulting construction provides an initiating ATG codon at the start of the glucoamylase signal, and an in frame ATG (Met) codon at the start of the TGF- $\alpha$  peptide.

#### 20 D. Expression and secretion of TGF- $\alpha$ in yeast cultures

Plasmid DNA (2-10  $\mu$ g/ml) was used to transform *S. cerevisiae* strain F762 (Mata, trp1 1, ura3-52) using standard techniques [Sherman et al., Methods in Yeast Genetics, Cold Spring Harbor, (1986)], selecting for tryptophan prototrophy. Strain F762 was obtained from Phillip Hunter, Department of Molecular Biology and Genetics, Johns Hopkins Medical School, Baltimore, Maryland. Cultures of selected transformants were grown in YNB (Difco) containing 2% glucose and uracil (20  $\mu$ g/ul). For plasmids utilizing the CUP1 promoter, copper sulfate at 100 micromolar was included. Cultures were harvested in late log or stationary phase. The cells were spun, and both the culture medium and cells retained. The cell pellets were washed, broken with glass beads in 0.05M acetate buffer, pH 5.0, and then the suspension centrifuged at 5000xg to remove the glass beads and cell debris. The soluble cell extracts and the culture fluids were then dialysed against 2% acetic acid overnight, using tubing of 2,000 molecular weight cut-off.

#### 35 E. Biological assay of TGF- $\alpha$

The dialysed cell lysates and culture fluids were analysed for TGF- $\alpha$  activity by three methods.

##### 40 (1) TGF- $\alpha$ /EGF Radioreceptor Assay

A431A human epidermoid carcinoma cells ( $1 \times 10^5$ ) were plated overnight in 24-well tissue culture dishes in DMEM containing 10% FCS at 37°C in 10% CO atmosphere. Cells were washed 3 times with binding buffer (Hank's balanced salts solution, 1 mg/ml bovine serum albumen (BSA) and 10 mM Hepes, pH 7.4) at 4°C. Binding buffer (100  $\mu$ l) containing 1 nM  $^{125}$ I TGF- $\alpha$  and various concentrations of the sample to be tested or epidermal growth factor (EGF) standard were added simultaneously and incubated at 4°C for 2 hours. Cold competition was performed with a 1000-fold excess of EGF. TGF- $\alpha$  was iodinated using the chloramine T procedure. Cells were washed with ice-cold binding buffer without BSA. The cells were solubilized for 30 min at 37°C with 0.5 ml of 0.5M NaOH. Radioactivity bound was determined using a gamma counter.

##### (2) Anchorage independent growth assay

Experimental samples which had been lyophilized, then dissolved in buffer containing 10% FCS and 0.05% BSA. The cells were added to an 0.8 ml bottom layer of DMEM containing 10% FCS and 0.05% BSA.

in 35 mm tissue culture dishes (Costar, Cambridge, Mass.). Plates were incubated for 8-10 days in 10% CO<sub>2</sub> atmosphere at 37°C. Colonies formed were stained with p-iodonitrotetrazolium violet overnight and counted on a Bausch and Lomb Image Analysis. System programmed to detect colonies greater than 60 µm in diameter.

### (3) Radioimmune assay

Assays were performed using a commercially available TGF-α RIA kit (Biotape), according to the manufacturers instructions.

### F. Activity in TGF-α produced in Yeast

The activity of TGF-α produced by plasmid pYSK151 and pYSK152 in shake flask cultures harvested 12 hours after reaching stationary phase was determined. The maximal yield obtained in the medium was approximately 2 mg/liter at OD<sub>540</sub> = 2.0. Greater than 95% of the total activity was found in the medium as compared to the cell extract.

TABLE 1

Estimated TGF-α Activity (ug/10 <sup>8</sup> cells)		
Plasmid	Medium	Cell extract
EGF competition binding assay:		
pYSK151	2.87	0.143
pYSK152	2.70	0.135
Soft Agar Assay:		
pYSK151	0.718	0.006
pYSK152	0.617	0.005
Radioimmune Assay:		
pYSK151	2.31	
pYSK152	2.24	

### G. Purification of TGF-α

Fermentation broth was centrifuged to remove cells, then filtered. The pH of the filtrate was adjusted to pH 2.7 using glacial acetic acid, then pumped through a column of Whatman Partisil ODS-3, (column dimensions: 48 x 650 mm packing 40-60 µm) at a rate of 200 ml/min which had been equilibrated to 0.1% trifluoroacetic acid (TFA). After loading, the column was washed with 2 column volumes of 0.1% TFA, and 5% i-propanol/0.1% TFA each. Crude TGF-α was eluted with 40% i-propanol/0.1% TFA.

Fractions were analyzed by SDS PAGE electrophoresis. Those fractions containing TGF-α were pooled together and lyophilized.

The powder (14 mg) was loaded onto a Vydac C<sub>18</sub> column (10 mm x 250 mm) using a gradient 16%

acetonitrile/0.05% TFA to 26% acetonitrile/0.05% TFA in 90 minutes. Fractions were collected and analyzed by HPLC. Recovery was approximately 30% of total activity. Three pools containing protein were made and were submitted for EGF/TGF radioreceptor assay, amino acid composition, and sequencing.

All three proteins were found to be biologically active, and varied in chain length at the N-terminal. The sequences were that of TGF- $\alpha$  (50%), Met-TGF- $\alpha$  (18%), and Gly-Ala-Met-TGF- $\alpha$  (22%). Comparison of amino acid composition and receptor binding assay showed that each had a specific activity of 1.0 EGF equivalents.

#### 10 EXAMPLE 12 - EXPRESSION AND SECRETION OF ILI- $\beta$ AND IL1-a

Human ILI- $\beta$  is expressed in a variety of cell types as a large (31 Kd) precursor polypeptide which is processed and secreted as a mature, unglycosylated, bioactive 17 Kd carboxyl-terminal fragment. Although ILI- $\beta$  is normally secreted, it lacks a signal sequence.

15 With the same methods of Example 11, the signal sequence derived from the cloned glucoamylase gene of *C. albicans* in combination with either the inducible CUP1 or constitutive TDH3 promoter in plasmids pYSK140 and pYSK142, was used to direct the synthesis and secretion of human recombinant interleukin-1  $\beta$  in *S. cerevisiae*. cDNA coding for the authentic mature form of ILI- $\beta$  (carried on a Ban1 fragment which was blunt-ended with Klenow polymerase) was fused as described in Example 11 to the  
20 glucoamylase signal sequence at the unique Sma1 site (see Fig. 4) of the pYSK140 and pYSK142, and transformed into yeast.

Each construct directs the efficient secretion of biologically active ILI- $\beta$ . Greater than 95% of the protein expressed is extracellular, resulting in accumulated levels of more than 2 mg/l as determined by Western blotting. Integration of a single copy of a TDH3 glucoamylase signal sequence ILI- $\beta$  expression cassette into  
25 the genome at the URA3 locus on chromosome V of *S. cerevisiae* resulted in a strain (GL46: MAT $\beta$  leu2-3,2-112 trp1-1 ade2-1 his3-11,3-15 can1-100 ura3 ILI- $\beta$ ) which secretes approximately 6 mg/l of product when grown to stationary phase in shake flasks under nonselective conditions (rich media). The amount of extracellular product obtained can be increased by constructing strains containing multiple integrations and/or by growing cells in fed-batch culture.

30 The secreted form of IL1- $\beta$  in yeast consists of approximately equimolar amounts of two protein species (17 Kd and 22 Kd). Pulse-chase experiments performed in the presence or absence of tunicamycin, followed by immunoprecipitation, reveal that the 22 Kd species is a glycosylated form of the molecule. A potential N-linked glycosylation site resides near the amino-terminus. The reason for the appearance of a mixture of two extracellular forms (glycosylated and unglycosylated) is unclear. Kinetic data indicate that the  
35 newly secreted product consists primarily of the 22 Kd glycosylated species which may then be converted, via either proteolytic cleavage or by an endoglycosidase, to the 17 Kd form. Direct fusion of the IL1B sequence to the CUP1 inducible promoter without a signal sequence results in expression of the 17 Kd intracellular form, plus a single 17 Kd extracellular species which accumulates in the medium of low phase cultures.

40 Purified extracellular IL1- $\beta$  is presently being sequenced to determine the primary signal peptide cleavage site as well as any secondary sites that may be recognized in yeast.

Using similar methods to those outlined in Example 11, the signal sequence derived from the cloned glucoamylase gene of *C. albicans*, was used to direct the synthesis and secretion of human recombinant interleukin-1  $\alpha$  in *S. cerevisiae*.

45

#### EXAMPLE 13 - EXPRESSION AND SECRETION OF B. PERTUSSIS TOXIN S1 SUBUNIT

50 Pertussis toxin, a promising protective antigen for potential use as a vaccine against whooping cough, is a hexameric protein composed of five different subunits. One of the subunits (S1) expresses an enzymatic activity, which is responsible for the toxicity of the protein. However S1 without the other subunits is devoid of toxic activities. The expression of the S1 gene and the secretion of the recombinant product should facilitate production and purification of this subunit, without contamination from the other pertussis toxin

peptide.

### A. Plasmid constructs and *S. cerevisiae* clones

A DNA fragment starting at the second amino acid following signal peptide cleavage site of S1 and extending to the natural TAG translational stop codon encoding an almost complete mature S1 protein was fused, in frame, to the signal peptide coding sequence of the *Candida albicans* glucoamylase gene. The fused coding sequences are preceded by the 5' non coding sequences of the glucoamylase gene including the promoter and transcription initiation sites of the STA gene (see Fig. 1). The expression cassette was transferred to an *E. coli*/*S. cerevisiae* shuttle vector containing the elements of the vector of Fig. 2 minus the env expression cassette, yielding plasmid pRIT13072. Plasmid pRIT13072 contains all the components of the vector of Fig. 2, except that the S1 expression cassette replaces the env expression cassette, and the ligation site on the right side of the cassette is an XbaI/Sal site, instead of an EspI/SalI site. pRIT13072 was transformed into a *S. cerevisiae* strain (genotype: *Mat $\alpha$  leu2 ura3 ade1 gln1*). A transformed yeast clone, 745.1 [pRIT13072], was selected for further work.

A Sau3a DNA restriction fragment encoding 186 amino acids of the S1 protein (amino acids 2 to 187 of the mature protein) was inserted into the coding sequence of the STA gene (see, Fig. 1). The insert was introduced at an engineered BamHI restriction site located near the carboxy terminal end of the glucoamylase protein. The resulting hybrid gene should express a fusion glucoamylase-S1 protein. The entire expression cassette is carried on an *E. coli*/*S. cerevisiae* shuttle plasmid vector similar to the one described above. The resulting plasmid, pRIT13073, was introduced into a *S. cerevisiae* strain (genotype: *Mat $\alpha$  leu2 ura3 ade1 gln1*) and a transformed clone, 745.1 [pRIT13073], was selected and used for subsequent analyses.

### B. Expression

The S1 subunit synthesized in clone 745.1 [pRIT13072] was readily detectable in Western blots of whole cell extracts using anti-pertussis toxin monoclonal antibody B2F8. The electrophoretic mobility of the recombinant protein in SDS-PAGE appeared to be similar to the mobility of the S1 subunit purified from *Bordetella pertussis* holotoxin, indicating that the glucoamylase signal peptide is removed and the S1 subunit processed in the yeast cell. Signal peptide cleavage usually occurs during or after translocation of the synthesized protein through the endoplasmic reticulum membrane. The processed S1 subunit should therefore be associated with the crude membrane fraction of yeast cell extracts and/or secreted into the culture medium. Yeast cell extracts were fractionated by differential centrifugation and the different fractions analyzed for the presence of S1. The S1 subunit was associated exclusively with the insoluble material after a 30000 xg centrifugation of the extracts. The pelleted fractions could be solubilized with 1% SDS or 4M urea. A number of other detergents [sodiumdeoxycholate (DOC, Sigma), Triton X100, 3-[(3-cholamidopropyl)-dimethyl ammonio]-1-propane sulfonate (CHAPS, Sigma) and zwittergen] were tested in an attempt to extract the S1 protein from yeast membrane. None had any significant effect on S1 solubilization.

Negative control and S1 containing cell extracts, before and after fractionation, were analysed for the presence of NAD glycohydrolase activities. None of the fractions contained significant enzymatic activities, but all of them strongly inhibited the NAD glycohydrolase activity of S1. No S1 protein could be detected by Western blot or by enzymatic activity analysis, in the culture medium.

Clone 745.1 [pRIT13073] was analyzed for the presence of a glucoamylase-S1 fusion protein (i.e., S1-GA protein). Glucoamylase enzymatic activity was tested by measuring the amount of glucose liberated from the substrate maltotriose at 37°C, in a reaction buffer containing limit NaPO<sub>4</sub>, 1% maltotriose, pH 6.5. S1 (NAD-glycohydrolase) activity was measured by the amount (pmoles) nicotinamide released per minute per ml of medium. The parent vector containing no expression cassette showed no detectable glucoamylase or S1 activity in culture medium or extracts. The parent vector containing just a glucoamylase expression cassette (i.e. the STA gene) showed glucoamylase activity in both culture medium and cell extracts, but no S1 activity. In contrast, the yeast strain carrying pRIT13073, the vector containing the glucoamylase S1 fusion expression cassette showed glucoamylase activity in both medium and extract and S1 activity in medium. Thus, *S. cerevisiae* cells transformed with plasmid pRIT13073 synthesize and

secrete the S1 subunit of *B. pertussis* toxin. The S1 subunit of *B. pertussis* toxin can be synthesized in yeast under the control of the glucoamylase or STA gene promoter, secreted into the yeast membranes via the glucoamylase signal peptide or exported into the culture medium as a fusion protein with glucoamylase. In

the latter case (i.e. the fusion glucoamylase-S1 protein) the S1 subunit is enzymatically active.

### C. Immunoprotective activity

Two mice were each injected with 35  $\mu$ g S1-GA(pRIT13073) (the S1-glucoamylase fusion protein prepared in Example 13, part B from the transformation of *S. cerevisiae* with pRIT13073) or glucoamylase (GA) in CFA, at day 0, and twice in IFA at days 30 and 58. The GA and S1-GA(pRIT13073) fusion protein used were both synthesized in *S. cerevisiae* and partially purified from the culture supernatant in a non-denaturing manner, such as by DEAE-cellulose chromatography, prior to the injection into mice. The mice were bled at days 54 and 75. The sera were analyzed by ELISA for the presence of anti-PT, anti-S1 and anti-rSld antibodies, and by the Chinese Hamster Ovary (CHO) Cell assay (Hewlett et al. (1983) Infect. Immun. 55, 24-38) for the presence of PT neutralizing antibodies. All mice injected with S1-GA(pRIT13073) fusion protein had increased anti-PT, anti-S1 and anti-rSld antibody titers. Antibody titers against these antigens did not significantly increase in the mice immunized with GA. All mice injected with S1-GA(pRIT13073) also had increased CHO cell neutralization titers. The arithmetic mean value was 920 for S1-GA(pRIT13073). These results show that the S1-GA protein is sufficient to induce a neutralizing and protective response, and that such protein is useful in a vaccine against pertussis.

In a manner similar to the method outlined in Example 13, parts A and B, the Sau3a DNA restriction fragments encoding 186 amino acids of the S1 protein (amino acids 2 to 187 of the mature protein) could be fused, in frame, to the signal peptide coding sequence of the Candida albicans glucoamylase gene, the resulting expression cassette could be transferred to an appropriate vector, the resulting vector could be transferred into an appropriate host S. cerevisiae strain, and the resulting protein could be used in a vaccine against pertussis.

### EXAMPLE 14-PARENTERAL VACCINE PREPARATION

5-25  $\mu$ g of S1-GA(pRIT13073), prepared according to the method of Example 13, is mixed with an aluminum adjuvant, such as aluminum hydroxide, to produce a vaccine in a form appropriate for incorporation into a parenteral administration dosage form.

While the above descriptions and examples fully describe the invention and the preferred embodiments thereof, it is understood that the invention is not limited to the particular disclosed embodiments.

This invention is not limited to the genes and promoters of Candida given in the examples above. Rather, this invention provides a general method for locating Candida gene expression units, promoters, regulatory regions and structural genes as well as a general method for expressing such functional DNA sequences in Saccharomyces.

Further, the heterologous genes expressed under control of the C. albicans glucoamylase signal sequence and yeast promoters are not limited to the exemplary proteins described herein. One of skill in the art will acknowledge the use of the described expression system for expressing and secreting a variety of heterologous genes of interest.

Thus, the invention includes all embodiments coming within the scope of the following claims.

### Claims

1. A transformed host microorganism of the genus Saccharomyces which comprises a functional DNA sequence derived from the same or substantially the same coding sequence as the signal sequence of the C. albicans glucoamylase gene.

2. The microorganism of Claim 1 wherein the functional DNA sequence comprises the same or substantially the same DNA sequence as that of Fig. 3.

3. The microorganism of Claim 1 wherein the functional DNA sequence comprises said signal sequence preceded by a yeast promoter.

4. The microorganism of Claim 1 wherein the functional DNA sequence comprises the same or substantially the same DNA sequence as that of Fig. 3, and wherein said signal sequence is fused in proper orientation with a yeast promoter.

5. The microorganism of Claim 1 wherein said signal sequence is fused in proper orientation with a yeast promoter.

selected heterologous gene.

7. The microorganism of Claim 6 wherein said heterologous gene is selected from the group consisting of TGF- $\alpha$ , Interleukin-1 $\beta$ , Interleukin-1 $\alpha$ , S1-GA protein, an HIV-I protein or fragments thereof.

8. The microorganism of Claim 1 which belongs to the species Saccharomyces cerevisiae.

5 9. The microorganism of Claim 1 which comprises a recombinant plasmid, wherein said plasmid comprises a replicator region and a functional DNA sequence derived from the same or substantially the same sequence as the signal sequence of the C. albicans glucoamylase gene.

10. A recombinant DNA plasmid comprising a functional DNA sequence containing a coding sequence the same or substantially the same as the signal sequence of the C. albicans glucoamylase gene.

10 11. The plasmid of Claim 10 wherein the functional DNA sequence comprises the same or substantially the same DNA sequence as that of Fig. 3.

12. The plasmid of Claim 10 wherein the functional DNA sequence comprises said signal sequence preceded by a yeast promoter.

15 13. The plasmid of Claim 10 which comprises a functional DNA sequence the same or substantially the same as the promoter sequence of the C. albicans glucoamylase gene.

14. The plasmid of Claim 13 wherein the functional DNA sequence comprises the same or substantially the same DNA sequence as that of Fig. 1.

15. The plasmid of Claim 10 wherein said signal sequence is fused in proper orientation with a selected heterologous gene.

20 16. The plasmid of Claim 15 wherein said heterologous gene is selected from the group consisting of TGF- $\alpha$ , Interleukin-1 $\beta$ , Interleukin-1 $\alpha$ , S1-GA protein, an HIV-I protein or fragments thereof.

17. A method of expressing a functional DNA sequence comprising a sequence the same or substantially the same as the signal or promoter sequence of the C. albicans glucoamylase gene which comprises transforming a host microorganism of the genus Saccharomyces with the functional DNA sequence, and culturing the transformed host under suitable conditions such that the functional DNA sequence is expressed.

18. The method of Claim 17 wherein said signal sequence is fused, in phase, and in proper orientation, with a selected heterologous gene.

30 19. The method of Claim 18 wherein said heterologous gene is selected from the group consisting of TGF- $\alpha$ , Interleukin-1 $\beta$ , Interleukin-1 $\alpha$ , S1-GA protein, an HIV-I protein or fragments thereof.

20. The method according to Claim 17 wherein the host microorganism belongs to the species Saccharomyces cerevisiae.

21. A promoter sequence for use in yeast expression vectors comprising the same or substantially the same sequence as that of Fig. 1.

35 22. A signal sequence for use in yeast expression vectors enabling the secretion of heterologous genes fused thereto comprising the same or substantially the same sequence of Fig. 3.

23. An HIV-I glycoprotein gp160 produced in unprocessed form by culturing a transformed Saccharomyces host cell wherein said host cell comprises a gp160 gene sequence fused in phase and in proper orientation to a yeast promoter and the signal sequence of the Candida albicans glucoamylase gene.

40 24. The glycoprotein of claim 23 produced in micellar form.

25. TGF- $\alpha$  produced by culturing a transformed Saccharomyces host cell wherein said host cell comprises a TGF- $\alpha$  gene sequence fused in phase and in proper orientation to a yeast promoter and the signal sequence of the Candida albicans glucoamylase gene.

45 26. S1-GA protein produced by culturing a transformed Saccharomyces host cell wherein said host cell comprises a S1-GA protein coding sequence fused in phase and in proper orientation (a) to a yeast promoter and (b) within the Candida albicans glucoamylase gene provided that such fusion includes the signal sequence of the Candida albicans glucoamylase gene.

27. The protein of Claim 26 which is S1-GA(pRIT13073).

50 28. S1-GA protein produced by culturing a transformed Saccharomyces host cell wherein said host cell comprises a S1-GA protein coding sequence fused in phase and in proper orientation to a yeast promoter and the signal sequence of the Candida albicans glucoamylase gene.

29. IL1- $\beta$  produced by culturing a transformed Saccharomyces host cell wherein said host cell

comprises a IL1- $\beta$  gene sequence fused in phase and in proper orientation to a yeast promoter and the signal sequence of the Candida albicans glucoamylase gene.

31. A vaccine for stimulating protection against whooping cough wherein such vaccine comprises an



immunoprotective and non-toxic quantity of S1-GA protein.

32. The vaccine of Claim 31 which comprises S1-GA(pRIT13073) protein.

33. A method for inoculating a human against subsequent whooping cough which comprises administering the vaccine of Claim 31 to such human.

5 34. S1-GA protein.

35. S1-GA protein coding sequence.

36. A recombinant DNA molecule comprising the S1-GA coding sequence of Claim 35.

37. A recombinant DNA vector comprising the S1-GA coding sequence of Claim 35.

38. A synthetic DNA molecule comprising the S1-GA coding sequence of Claim 35.

10 39. A host cell transformed with the S1-GA coding sequence containing vector of Claim 37.

15

20

25

30

35

40

45

50

**FIG. 1**

TaqI

CGAGGATTATTGTTATTGTTGTTTCGTACGGAGATTTCTCCCTAAATTACACAAAGTACA  
 TCCTAATAACAATAACAACAAGCATGCCTCTAAAGAGGGATTTAATGTGTTTCATGT

ATAATTCCCCTTTCTTGTTACAAATTAGTGTACGTATGTAGCACCTGATTAGTTTATTTA  
 TATTAAGGGGAAAGAACAATGTTTAATCACATGCATACATCGTGGACTAATCAAATAAAT

TTTATTCATAGTCAGGCCAGAAAGTAATTACCAAGTTTAGTATTTCCGACCGCCCCCAA  
 AAATAAGTATCAGTCCGGTCTTTCATTAATGGTTCAAATCATAAAGGCTGGCGGGGGGTT

ACAGGAACCTTATTCCGAGCAAAAGTAGTTTATCTCCTGTACGAAAAGAAGATGTAATTAT  
 TGTCTTGAATAAGGCTCGTTTTTCATCAAATAGAGGACATGCTTTTCTTCTACATTAATA

TGCTACTAAAAACAGACAATATAAAAGCCCCGAAAAAGTAATTGCAAAATTCGTTTGT  
 ACGATGATTTTTTGTCTGTTATATTTTCGGGGGCTTTTTTCATTAACGTTTAAAGCAAACA

TAAGAATTTTCAAACCAAGACAAACCAACCAACCAACCAACCAAGAAAATATGAAA  
 ATTCTTAAAAGTTTGTTCTGTTTGGTTTGGTTTGGTTTGGTTTGGTTCTTTTATACTTT  
 MetLys

Fnu4H1

TTATTATCAAATTCATTGTTACTGCTTTGGGTCTTACTTCAATTGTCAATGC  
 AATAATAGTTTTAAGTAACAATGACGAAACCCAGAATGAAGTTAACAGTTACGG  
 LeuLeuSerLysPheIleValThrAlaLeuGlyLeuThrSerIleValAsnAla

**FIG. 3**

AAACCAAGAAAATATGAAATTATTATCAAATTCATTGTTACTGCTTTGGGTCTTACTTCA  
 METLysLeuLeuSerLysPheIleValThrAlaLeuGlyLeuThrSer

ATTGTCAATGCCGCTCCAACAAGTAGTAGTAGT  
 IleValAsnAlaAlaProThrSerSerSerSer

|  
 processing site




FIG. 2

## FIG. 4

XhoI HpaI  
 \* \*  
 TCGAGAATGAAATTATTATCAAAATTCATTGTTACTGCTTTGGGTCTTACTTCAATTGTTAAC  
 CTTACTTTAATAATAGTTTTTAAGTAACAATGACGAAACCCAGAATGAAGTTAACAATTG  
 METLysLeuLeuSerLysPheIleValThrAlaLeuGlyLeuThrSerIleValAsn

SmaI NcoI  
 \*  
 GCCGCTCCCGGGGC  
 CGGCGAGGGCCCCGGTAC  
 AlaAlaProGlyAlaMet

## FIG. 5

Hind III NcoI DdeI  
 .  
 AAGCTTTTCCATGGTGGTGTCCCATTTTAATGACTGCCCAGATTCCCACACTCAGTTCTGC  
 TTCGAAAAGGTACCACCACAGGGTAAAATTACTGACGGOTCTAAGGGTGTGAGTCAAGACG  
 SerPheSerMetValValSerHisPheAsnAspCysProAspSerHisThrGlnPheCys

PstI SphI  
 C . T  
 TTTCATGGAACCTGCAGGTTTTTGGTGCAGGAGGACAAGCCAGCATGCGTCTGCCATTCT  
 AAAGTACCTTGGACGTCCAAAACCACGTCCTCCTGTTTCGGTCGTACGCAGACGGTAAGA  
 PheHisGlyThrCysArgPheLeuValGlnGluAspLysProAlaCysValCysHisSer

Sali  
 AccI  
 HincII  
 DraIII  
 . G C ...  
 GGGTACGTTGGTGCACGCTGTGAACATGCGGATCTCCTGGCTTAAGTCGAC  
 CCCATGCAACCACGTGCCACACTGTACGCCTAGAGGACCGAATTCAGCTG  
 GlyTyrValGlyAlaArgCysGluHisAlaAspLeuLeuAlaEndValAsp